



# Myosin IIa (phospho-Ser628) rabbit pAb

<b>Catalog No</b>	YP-Ab-17220
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat
<b>Applications</b>	WB;IHC
<b>Gene Name</b>	MYH9
<b>Protein Name</b>	Myosin IIa (Ser1943)
<b>Immunogen</b>	Synthesized phosho peptide around human Myosin IIa (Ser1943)
<b>Specificity</b>	This antibody detects endogenous levels of Human Mouse Rat Myosin IIa (phospho-Ser1943)
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Dilution</b>	WB 1:500-2000;IHC-p 1:50-300
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	Myosin-9 (Cellular myosin heavy chain, type A) (Myosin heavy chain 9) (Myosin heavy chain, non-muscle IIa) (Non-muscle myosin heavy chain A) (NMMHC-A) (Non-muscle myosin heavy chain IIa) (NMMHC II-a) (NMMHC-IIA)
<b>Observed Band</b>	215kD
<b>Cell Pathway</b>	Cytoplasm, cytoskeleton . Cytoplasm, cell cortex . Cytoplasmic vesicle, secretory vesicle, Cortical granule . Colocalizes with actin filaments at lamellipodia margins and at the leading edge of migrating cells (PubMed:20052411). In retinal pigment epithelial cells, predominantly localized to stress fiber-like structures with some localization to cytoplasmic puncta (PubMed:27331610). .
<b>Tissue Specificity</b>	In the kidney, expressed in the glomeruli. Also expressed in leukocytes.
<b>Function</b>	disease:Defects in MYH9 are the cause of Alport syndrome with macrothrombocytopenia (APSM) [MIM:153650]. APSM is an autosomal dominant disorder characterized by the association of ocular lesions, sensorineural hearing loss and nephritis (Alport syndrome) with platelet defects..disease:Defects in MYH9 are the cause of Epstein syndrome (EPS) [MIM:153650]. EPS is an autosomal dominant disorder characterized by the association of macrothrombocytopathy, sensorineural hearing loss and nephritis..disease:Defects in MYH9 are the cause of Fechtner syndrome (FTNS) [MIM:153640]. FTNS is an autosomal dominant macrothrombocytopenia characterized by thrombocytopenia, giant platelets and leukocyte inclusions that



are small and poorly organized. Additionally, FTNS is distinguished by Alport-like clinical features of sensorineural deafness, cataracts and nephritis.,disease:Defects in MYH9 are the cause o

#### Background

This gene encodes a conventional non-muscle myosin; this protein should not be confused with the unconventional myosin-9a or 9b (MYO9A or MYO9B). The encoded protein is a myosin IIA heavy chain that contains an IQ domain and a myosin head-like domain which is involved in several important functions, including cytokinesis, cell motility and maintenance of cell shape. Defects in this gene have been associated with non-syndromic sensorineural deafness autosomal dominant type 17, Epstein syndrome, Alport syndrome with macrothrombocytopenia, Sebastian syndrome, Fechtner syndrome and macrothrombocytopenia with progressive sensorineural deafness. [provided by RefSeq, Dec 2011],


#### matters needing attention

Avoid repeated freezing and thawing!

#### Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

### Products Images



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).