

(Tel: 400-999-8863 **(** Emall:Upingbio.163.com





PTPRR rabbit pAb

Catalog No YP-Ab-17251 Isotype IgG Reactivity Human, Mouse,Rat Applications IHC,WB Gene Name PTPRR ECPTP PTPRQ Protein Name Receptor-type tyrosine-protein phosphatase R (R-PTP-R) (EC 3.1.3.48) (Ch-PTPase) (NC-PTPCOM1) (Protein-tyrosine phosphatase PCPTP1) Immunogen Synthesized peptide derived from human N-ternal PTPRR Specificity This antibody detects endogenous levels of PTPRR at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Rabbit,polyclonal Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 IHC 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Receptor-type tyrosine-protein phosphatase R (R-PTP-R) (EC 3.1.3.48) (Ch-1PTPase) (NC-PTPCOM1) (Protein-tyrosine phosphatase PCPTP1) Observed Band Cell Pathway Isoform Alphaj: Cell membrane; Single-pass type I membrane protein.; Ilsoform beltaj: Cytoplasm, perinuclear reas within the cytoplasm. Distaje: Cytoplasm, perinuclear reas within the cytoplasm. Exp		
Reactivity Human, Mouse,Rat Applications IHC,WB Gene Name PTPRR ECPTP PTPRQ Protein Name Receptor-type tyrosine-protein phosphatase R (R-PTP-R) (EC 3.1.3.48) (Ch-1PTPase) (NC-PTPCOM1) (Protein-tyrosine phosphatase PCPTP1) Immunogen Synthesized peptide derived from human N-ternal PTPRR Specificity This antibody detects endogenous levels of PTPRR at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Rabbit,polyclonal Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 IHC 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Receptor-type tyrosine-protein phosphatase R (R-PTP-R) (EC 3.1.3.48) (Ch-1PTPase) (NC-PTPCOM1) (Protein-tyrosine phosphatase PCPTP1) Observed Band Cell Pathway [Isoform Alpha]: Cell membrane; Single-pass type I membrane protein.: [Isoform Delta]: Cytoplasm, perinuclear region. Locates to the perinuclear areas within the cytoplasm. Tissue Specificity Expressed in brain, placenta, small intestine, stomach, uterus and weakly in the prostate. Isoform alpha has been observed only in the brain. Isoform gamma is expressed in brain, placenta, prostate, small intestine, and uterus. Sequesters mitogen-activated protein kinases (MAPKs) such as MAPK1, MAPK3 and MAPK14 in the cytoplasm in an inactive form. The MAPKs bind to a dephosphorylated kinase interacting motif, phosphorylation of which by the protein kinase A complex releases the MAPKs for activation and translocation into the nucleus (By similarity).	Catalog No	YP-Ab-17251
Applications IHC,WB Gene Name PTPRR ECPTP PTPRQ Protein Name Receptor-type tyrosine-protein phosphatase R (R-PTP-R) (EC 3.1.3.48) (Ch-1PTPase) (NC-PTPCOM1) (Protein-tyrosine phosphatase PCPTP1) Immunogen Synthesized peptide derived from human N-ternal PTPRR Specificity This antibody detects endogenous levels of PTPRR at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Rabbit,polyclonal Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 IHC 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Receptor-type tyrosine-protein phosphatase R (R-PTP-R) (EC 3.1.3.48) (Ch-1PTPase) (NC-PTPCOM1) (Protein-tyrosine phosphatase PCPTP1) Observed Band Cell Pathway [Isoform Alpha]: Cell membrane; Single-pass type I membrane protein.; [Isoform Delta]: Cytoplasm, perinuclear region. Locates to the perinuclear areas within the cytoplasm. Tissue Specificity Expressed in brain, placenta, small intestine, stomach, uterus and weakly in the prostate. Isoform alpha has been observed only in the brain. Isoform gamma is expressed in brain, placenta, prostate, small intestine and uterus. Sequesters mitogen-activated protein kinases (MAPKs) such as MAPK1, MAPK3 and MAPK14 in the cytoplasm in an inactive form. The MAPKs bind to a dephosphorylated kinase interacting motif, phosphorylation of which by the protein kinase A complex releases the MAPKs for activation and translocation into the nucleus (By similarity).	Isotype	IgG
Gene Name PTPRR ECPTP PTPRQ Protein Name (Receptor-type tyrosine-protein phosphatase R (R-PTP-R) (EC 3.1.3.48) (Ch-1PTPase) (NC-PTPCOM1) (Protein-tyrosine phosphatase PCPTP1) Immunogen Synthesized peptide derived from human N-ternal PTPRR Specificity This antibody detects endogenous levels of PTPRR at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Rabbit,polyclonal Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 IHC 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Receptor-type tyrosine-protein phosphatase R (R-PTP-R) (EC 3.1.3.48) (Ch-1PTPase) (NC-PTPCOM1) (Protein-tyrosine phosphatase PCPTP1) Observed Band Cell Pathway [Isoform Alpha]: Cell membrane; Single-pass type I membrane protein; [Isoform Delta]: Cytoplasm; [Isoform Gamma]: Cytoplasm, perinuclear region. Locates to the perinuclear areas within the cytoplasm; [Isoform Gamma]: Cytoplasm, perinuclear region. Locates to the prostate. Isoform alpha has been observed only in the brain. Isoform gamma is expressed in brain, placenta, small intestine, stomach, uterus and weakly in the prostate. Isoform alpha has been observed only in the brain. Isoform gamma is expressed in brain, placenta, placenta, mall intestine and uterus. Soform delta is expressed in brain, placenta, placenta in inactive form. The MAPKs bind to a dephosphorylated kinase interacting motif, phosphorylation of which by the protein kinase A complex releases the MAPKs for activation and translocation into the nucleus (By similarity).	Reactivity	Human, Mouse,Rat
Protein Name Receptor-type tyrosine-protein phosphatase R (R-PTP-R) (EC 3.1.3.48) (Ch-1PTPase) (NC-PTPCOM1) (Protein-tyrosine phosphatase PCPTP1) Immunogen Synthesized peptide derived from human N-ternal PTPRR Specificity This antibody detects endogenous levels of PTPRR at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Rabbit,polyclonal Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 IHC 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Receptor-type tyrosine-protein phosphatase R (R-PTP-R) (EC 3.1.3.48) (Ch-1PTPase) (NC-PTPCOM1) (Protein-tyrosine phosphatase PCPTP1) Observed Band Cell Pathway [Isoform Alpha]: Cell membrane; Single-pass type I membrane protein; [Isoform Delta]: Cytoplasm; [Isoform Gamma]: Cytoplasm, perinuclear region. Locates to the perinuclear areas within the cytoplasm; [Isoform Gamma]: Cytoplasm, perinuclear region. Locates to the perinuclear areas within the cytoplasm, placenta, small intestine, stomach, uterus and weakly in the prostate. Isoform alpha has been observed only in the brain. Isoform gamma is expressed in brain, placenta, placenta, placenta, placenta, placenta and uterus. Isoform delta is expressed in brain, placenta, placenta and uterus. Seguesters mitogen-activated protein kinases (MAPKs) such as MAPK1, MAPK3 and MAPK14 in the cytoplasm in inactive form. The MAPKs bind to a dephosphorylated kinase interacting motif, phosphorylation of which by the protein kinase A complex releases the MAPKs for activation and translocation into the nucleus (By similarity).	Applications	IHC,WB
Immunogen Synthesized peptide derived from human N-ternal PTPRR	Gene Name	PTPRR ECPTP PTPRQ
Specificity This antibody detects endogenous levels of PTPRR at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Rabbit,polyclonal Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 IHC 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Receptor-type tyrosine-protein phosphatase R (R-PTP-R) (EC 3.1.3.48) (Ch-1PTPase) (NC-PTPCOM1) (Protein-tyrosine phosphatase PCPTP1) Observed Band Cell Pathway [Isoform Alpha]: Cell membrane; Single-pass type I membrane protein.; [Isoform Delta]: Cytoplasm, perinuclear region. Locates to the perinuclear areas within the cytoplasm.; Isoform Gamma]: Cytoplasm, perinuclear region. Locates to the perinuclear areas within the cytoplasm. Tissue Specificity Expressed in brain, placenta, small intestine, stomach, uterus and weakly in the prostate. Isoform alpha has been observed only in the brain. Isoform gamma is expressed in brain, placenta and uterus. Isoform delta is expressed in brain, kidney, placenta, prostate, small intestine and uterus. Function Sequesters mitogen-activated protein kinases (MAPKs) such as MAPK1, MAPK3 and MAPK14 in the cytoplasm in an inactive form. The MAPKs bind to a dephosphorylated kinase interacting motif, phosphorylation of which by the protein kinases complex releases the MAPKs for activation and translocation into the nucleus (By similarity).	Protein Name	Receptor-type tyrosine-protein phosphatase R (R-PTP-R) (EC 3.1.3.48) (Ch-1PTPase) (NC-PTPCOM1) (Protein-tyrosine phosphatase PCPTP1)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Rabbit,polyclonal Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 IHC 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Receptor-type tyrosine-protein phosphatase R (R-PTP-R) (EC 3.1.3.48) (Ch-1PTPase) (NC-PTPCOM1) (Protein-tyrosine phosphatase PCPTP1) Observed Band Cell Pathway [Isoform Alpha]: Cell membrane; Single-pass type I membrane protein.; [Isoform Delta]: Cytoplasm, perinuclear region. Locates to the perinuclear areas within the cytoplasm.] Isoform Gamma]: Cytoplasm, perinuclear region. Locates to the perinuclear areas within the cytoplasm. Expressed in brain, placenta, small intestine, stomach, uterus and weakly in the prostate. Isoform alpha has been observed only in the brain. Isoform gamma is expressed in brain, placenta, prostate, small intestine and uterus. Function Sequesters mitogen-activated protein kinases (MAPKs) such as MAPK1, MAPK3 and MAPK14 in the cytoplasm in an inactive form. The MAPKs bind to a dephosphorylated kinase interacting motif, phosphorylation of which by the protein kinase A complex releases the MAPKs for activation and translocation into the nucleus (By similarity).	Immunogen	Synthesized peptide derived from human N-ternal PTPRR
Source Rabbit,polyclonal Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 IHC 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Receptor-type tyrosine-protein phosphatase R (R-PTP-R) (EC 3.1.3.48) (Ch-1PTPase) (NC-PTPCOM1) (Protein-tyrosine phosphatase PCPTP1) Observed Band Cell Pathway [Isoform Alpha]: Cell membrane; Single-pass type I membrane protein.; [Isoform Delta]: Cytoplasm., perinuclear region. Locates to the perinuclear areas within the cytoplasm.; Isoform Gamma]: Cytoplasm, perinuclear region. Locates to the perinuclear areas within the cytoplasm. Tissue Specificity Expressed in brain, placenta, small intestine, stomach, uterus and weakly in the prostate. Isoform alpha has been observed only in the brain. Isoform gamma is expressed in brain, placenta and uterus. Isoform delta is expressed in brain, kidney, placenta, prostate, small intestine and uterus. Function Sequesters mitogen-activated protein kinases (MAPKs) such as MAPK1, MAPK3 and MAPK14 in the cytoplasm in an inactive form. The MAPKs bind to a dephosphorylated kinase interacting motif, phosphorylation of which by the protein kinase A complex releases the MAPKs for activation and translocation into the nucleus (By similarity).	Specificity	This antibody detects endogenous levels of PTPRR at Human, Mouse,Rat
Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 IHC 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Receptor-type tyrosine-protein phosphatase R (R-PTP-R) (EC 3.1.3.48) (Ch-1PTPase) (NC-PTPCOM1) (Protein-tyrosine phosphatase PCPTP1) Observed Band Cell Pathway [Isoform Alpha]: Cell membrane; Single-pass type I membrane protein.; [Isoform Delta]: Cytoplasm, perinuclear region. Locates to the perinuclear areas within the cytoplasm.; Isoform Gamma]: Cytoplasm, perinuclear region. Locates to the perinuclear areas within the cytoplasm. Expressed in brain, placenta, small intestine, stomach, uterus and weakly in the prostate. Isoform alpha has been observed only in the brain. Isoform gamma is expressed in brain, placenta and uterus. Isoform delta is expressed in brain, placenta, prostate, small intestine and uterus. Function Sequesters mitogen-activated protein kinases (MAPKs) such as MAPK1, MAPK3 and MAPK14 in the cytoplasm in an inactive form. The MAPKs bind to a dephosphorylated kinase interacting motif, phosphorylation of which by the protein kinase A complex releases the MAPKs for activation and translocation into the nucleus (By similarity).	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Dilution WB 1:500-2000 IHC 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Receptor-type tyrosine-protein phosphatase R (R-PTP-R) (EC 3.1.3.48) (Ch-1PTPase) (NC-PTPCOM1) (Protein-tyrosine phosphatase PCPTP1) Observed Band Cell Pathway [Isoform Alpha]: Cell membrane; Single-pass type I membrane protein.; [Isoform Delta]: Cytoplasm, perinuclear region. Locates to the perinuclear areas within the cytoplasm.; [Isoform Gamma]: Cytoplasm, perinuclear region. Locates to the perinuclear areas within the cytoplasm. Tissue Specificity Expressed in brain, placenta, small intestine, stomach, uterus and weakly in the prostate. Isoform alpha has been observed only in the brain. Isoform gamma is expressed in brain, placenta and uterus. Isoform delta is expressed in brain, kidney, placenta, prostate, small intestine and uterus. Function Sequesters mitogen-activated protein kinases (MAPKs) such as MAPK1, MAPK3 and MAPK14 in the cytoplasm in an inactive form. The MAPKs bind to a dephosphorylated kinase interacting motif, phosphorylation of which by the protein kinase A complex releases the MAPKs for activation and translocation into the nucleus (By similarity).	Source	Rabbit,polyclonal
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Receptor-type tyrosine-protein phosphatase R (R-PTP-R) (EC 3.1.3.48) (Ch-1PTPase) (NC-PTPCOM1) (Protein-tyrosine phosphatase PCPTP1) Observed Band Cell Pathway [Isoform Alpha]: Cell membrane; Single-pass type I membrane protein.; [Isoform Delta]: Cytoplasm, perinuclear region. Locates to the perinuclear areas within the cytoplasm.; [Isoform Gamma]: Cytoplasm, perinuclear region. Locates to the perinuclear areas within the cytoplasm. Tissue Specificity Expressed in brain, placenta, small intestine, stomach, uterus and weakly in the prostate. Isoform alpha has been observed only in the brain. Isoform gamma is expressed in brain, placenta and uterus. Isoform delta is expressed in brain, kidney, placenta, prostate, small intestine and uterus. Function Sequesters mitogen-activated protein kinases (MAPKs) such as MAPK1, MAPK3 and MAPK14 in the cytoplasm in an inactive form. The MAPKs bind to a dephosphorylated kinase interacting motif, phosphorylation of which by the protein kinase A complex releases the MAPKs for activation and translocation into the nucleus (By similarity).	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Receptor-type tyrosine-protein phosphatase R (R-PTP-R) (EC 3.1.3.48) (Ch-1PTPase) (NC-PTPCOM1) (Protein-tyrosine phosphatase PCPTP1) Observed Band Cell Pathway [Isoform Alpha]: Cell membrane; Single-pass type I membrane protein.; [Isoform Delta]: Cytoplasm, perinuclear region. Locates to the perinuclear areas within the cytoplasm.; [Isoform Gamma]: Cytoplasm, perinuclear region. Locates to the perinuclear areas within the cytoplasm. Tissue Specificity Expressed in brain, placenta, small intestine, stomach, uterus and weakly in the prostate. Isoform alpha has been observed only in the brain. Isoform gamma is expressed in brain, placenta and uterus. Isoform delta is expressed in brain, kidney, placenta, prostate, small intestine and uterus. Function Sequesters mitogen-activated protein kinases (MAPKs) such as MAPK1, MAPK3 and MAPK14 in the cytoplasm in an inactive form. The MAPKs bind to a dephosphorylated kinase interacting motif, phosphorylation of which by the protein kinase A complex releases the MAPKs for activation and translocation into the nucleus (By similarity).	Dilution	WB 1:500-2000 IHC 1:50-200
Synonyms Receptor-type tyrosine-protein phosphatase R (R-PTP-R) (EC 3.1.3.48) (Ch-1PTPase) (NC-PTPCOM1) (Protein-tyrosine phosphatase PCPTP1) Observed Band Cell Pathway [Isoform Alpha]: Cell membrane; Single-pass type I membrane protein.; [Isoform Delta]: Cytoplasm, perinuclear region. Locates to the perinuclear areas within the cytoplasm.; [Isoform Gamma]: Cytoplasm, perinuclear region. Locates to the perinuclear areas within the cytoplasm. Tissue Specificity Expressed in brain, placenta, small intestine, stomach, uterus and weakly in the prostate. Isoform alpha has been observed only in the brain. Isoform gamma is expressed in brain, placenta and uterus. Isoform delta is expressed in brain, kidney, placenta, prostate, small intestine and uterus. Function Sequesters mitogen-activated protein kinases (MAPKs) such as MAPK1, MAPK3 and MAPK14 in the cytoplasm in an inactive form. The MAPKs bind to a dephosphorylated kinase interacting motif, phosphorylation of which by the protein kinase A complex releases the MAPKs for activation and translocation into the nucleus (By similarity).	Concentration	1 mg/ml
Synonyms Receptor-type tyrosine-protein phosphatase R (R-PTP-R) (EC 3.1.3.48) (Ch-1PTPase) (NC-PTPCOM1) (Protein-tyrosine phosphatase PCPTP1) Observed Band Cell Pathway [Isoform Alpha]: Cell membrane; Single-pass type I membrane protein.; [Isoform Delta]: Cytoplasm, perinuclear region. Locates to the perinuclear areas within the cytoplasm.; [Isoform Gamma]: Cytoplasm, perinuclear region. Locates to the perinuclear areas within the cytoplasm. Tissue Specificity Expressed in brain, placenta, small intestine, stomach, uterus and weakly in the prostate. Isoform alpha has been observed only in the brain. Isoform gamma is expressed in brain, placenta and uterus. Isoform delta is expressed in brain, kidney, placenta, prostate, small intestine and uterus. Sequesters mitogen-activated protein kinases (MAPKs) such as MAPK1, MAPK3 and MAPK14 in the cytoplasm in an inactive form. The MAPKs bind to a dephosphorylated kinase interacting motif, phosphorylation of which by the protein kinase A complex releases the MAPKs for activation and translocation into the nucleus (By similarity).	Purity	≥90%
(Ch-1PTPase) (NC-PTPCOM1) (Protein-tyrosine phosphatase PCPTP1) Observed Band Cell Pathway [Isoform Alpha]: Cell membrane; Single-pass type I membrane protein.; [Isoform Delta]: Cytoplasm, perinuclear region. Locates to the perinuclear areas within the cytoplasm.; [Isoform Gamma]: Cytoplasm, perinuclear region. Locates to the perinuclear areas within the cytoplasm. Tissue Specificity Expressed in brain, placenta, small intestine, stomach, uterus and weakly in the prostate. Isoform alpha has been observed only in the brain. Isoform gamma is expressed in brain, placenta and uterus. Isoform delta is expressed in brain, kidney, placenta, prostate, small intestine and uterus. Function Sequesters mitogen-activated protein kinases (MAPKs) such as MAPK1, MAPK3 and MAPK14 in the cytoplasm in an inactive form. The MAPKs bind to a dephosphorylated kinase interacting motif, phosphorylation of which by the protein kinase A complex releases the MAPKs for activation and translocation into the nucleus (By similarity).	Storage Stability	-20°C/1 year
Cell Pathway [Isoform Alpha]: Cell membrane; Single-pass type I membrane protein.; [Isoform Delta]: Cytoplasm, perinuclear region. Locates to the perinuclear areas within the cytoplasm.; [Isoform Gamma]: Cytoplasm, perinuclear region. Locates to the perinuclear areas within the cytoplasm. Tissue Specificity Expressed in brain, placenta, small intestine, stomach, uterus and weakly in the prostate. Isoform alpha has been observed only in the brain. Isoform gamma is expressed in brain, placenta and uterus. Isoform delta is expressed in brain, kidney, placenta, prostate, small intestine and uterus. Function Sequesters mitogen-activated protein kinases (MAPKs) such as MAPK1, MAPK3 and MAPK14 in the cytoplasm in an inactive form. The MAPKs bind to a dephosphorylated kinase interacting motif, phosphorylation of which by the protein kinase A complex releases the MAPKs for activation and translocation into the nucleus (By similarity).	Synonyms	
Delta]: Cytoplasm, perinuclear region. Locates to the perinuclear areas within the cytoplasm.; [Isoform Gamma]: Cytoplasm, perinuclear region. Locates to the perinuclear areas within the cytoplasm. Tissue Specificity Expressed in brain, placenta, small intestine, stomach, uterus and weakly in the prostate. Isoform alpha has been observed only in the brain. Isoform gamma is expressed in brain, placenta and uterus. Isoform delta is expressed in brain, kidney, placenta, prostate, small intestine and uterus. Function Sequesters mitogen-activated protein kinases (MAPKs) such as MAPK1, MAPK3 and MAPK14 in the cytoplasm in an inactive form. The MAPKs bind to a dephosphorylated kinase interacting motif, phosphorylation of which by the protein kinase A complex releases the MAPKs for activation and translocation into the nucleus (By similarity).	Observed Band	
prostate. Isoform alpha has been observed only in the brain. Isoform gamma is expressed in brain, placenta and uterus. Isoform delta is expressed in brain, kidney, placenta, prostate, small intestine and uterus. Function Sequesters mitogen-activated protein kinases (MAPKs) such as MAPK1, MAPK3 and MAPK14 in the cytoplasm in an inactive form. The MAPKs bind to a dephosphorylated kinase interacting motif, phosphorylation of which by the protein kinase A complex releases the MAPKs for activation and translocation into the nucleus (By similarity).	Cell Pathway	Delta]: Cytoplasm, perinuclear region. Locates to the perinuclear areas within the cytoplasm.; [Isoform Gamma]: Cytoplasm, perinuclear region. Locates to the
MAPK3 and MAPK14 in the cytoplasm in an inactive form. The MAPKs bind to a dephosphorylated kinase interacting motif, phosphorylation of which by the protein kinase A complex releases the MAPKs for activation and translocation into the nucleus (By similarity).	Tissue Specificity	prostate. Isoform alpha has been observed only in the brain. Isoform gamma is expressed in brain, placenta and uterus. Isoform delta is expressed in brain,
Background	Function	MAPK3 and MAPK14 in the cytoplasm in an inactive form. The MAPKs bind to a dephosphorylated kinase interacting motif, phosphorylation of which by the protein kinase A complex releases the MAPKs for activation and translocation into
	Background	



UpingBio technology Co.,Ltd

(Tel: 400-999-8863 ■ Emall:Upingbio.163.com



matters	needing	
attention		

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images