



MBOA5 rabbit pAb

Catalog No	YP-Ab-17255
Isotype	IgG
Reactivity	Human, Mouse,Rat
Applications	IHC,WB
Gene Name	LPCAT3 MBOAT5 OACT5
Protein Name	Lysophospholipid acyltransferase 5 (LPLAT 5) (EC 2.3.1.-) (1-acylglycerophosphocholine O-acyltransferase) (EC 2.3.1.23) (1-acylglycerophosphoserine O-acyltransferase) (EC 2.3.1.n6) (Lysophosphatidylch
Immunogen	Synthesized peptide derived from human C-ternal MBOA5
Specificity	This antibody detects endogenous levels of MBOA5 at Human, Mouse,Rat
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Rabbit,polyclonal
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000 IHC 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Lysophospholipid acyltransferase 5 (LPLAT 5) (EC 2.3.1.-) (1-acylglycerophosphocholine O-acyltransferase) (EC 2.3.1.23) (1-acylglycerophosphoserine O-acyltransferase) (EC 2.3.1.n6) (Lysophosphatidylcholine acyltransferase) (LPCAT) (Lyso-PC acyltransferase) (Lysophosphatidylcholine acyltransferase 3) (Lyso-PC acyltransferase 3) (Lysophosphatidylserine acyltransferase) (LPSAT) (Lyso-PS acyltransferase) (Membrane-bound O-acyltransferase domain-containing protein 5) (O-acyltransferase domain-containing protein 5)
Observed Band	
Cell Pathway	Endoplasmic reticulum membrane ; Multi-pass membrane protein .
Tissue Specificity	Highly expressed in liver, pancreas and adipose tissue. Very low expression in skeletal muscle and heart. Detected in neutrophils.
Function	Lysophospholipid O-acyltransferase (LPLAT) that catalyzes the reacylation step of the phospholipid remodeling process also known as the Lands cycle . Catalyzes transfer of the fatty acyl chain from fatty acyl-CoA to 1-acyl lysophospholipid to form various classes of phospholipids. Converts 1-acyl



lysophosphatidylcholine (LPC) into phosphatidylcholine (PC) (LPCAT activity), 1-acyl lysophosphatidylserine (LPS) into phosphatidylserine (PS) (LPSAT activity) and 1-acyl lysophosphatidylethanolamine (LPE) into phosphatidylethanolamine (PE) (LPEAT activity) . Favors polyunsaturated fatty acyl-CoAs as acyl donors compared to saturated fatty acyl-CoAs . Has higher activity for LPC acyl acceptors compared to LPEs and LPSS. Can also transfer the fatty acyl chain from fatty acyl-CoA to 1-O-alkyl lysophospholipid or 1-O-alkenyl lysophospholipid with lower efficiency (By similarity). Acts as a major L

Background

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images