



# STING Rabbit pAb

<b>Catalog No</b>	YP-Ab-17777
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human
<b>Applications</b>	WB, ICC/IF, FC
<b>Gene Name</b>	STING1
<b>Alternative Names</b>	ERIS; MITA; MPYS; SAVI; NET23; STING; hMITA; hSTING; STING-beta; Transmembrane Protein 173
<b>Research Field</b>	Immunology
<b>Product Categories</b>	Primary antibody
<b>Host</b>	Rabbit
<b>Molecular Weight</b>	Calculated MW: 42 kDa; Observed MW: 37 kDa
<b>Clonality</b>	Polyclonal Antibody
<b>Clonality No.</b>	-
<b>Dilution</b>	WB: 1/500-1/1000 IF: 1/50-1/200 FC: 1/50-1/100
<b>Immunogen</b>	A synthesized peptide derived from human TMEM173
<b>Purification</b>	Affinity Chromatography
<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Form</b>	Liquid
<b>Buffer System</b>	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Background</b>	Facilitator of innate immune signaling that promotes the production of type I interferon (IFN-alpha and IFN-beta). Innate immune response is triggered in response to non-CpG double-stranded DNA from viruses and bacteria delivered to the cytoplasm.
<b>matters needing attention</b>	Avoid repeated freezing and thawing!



### Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

### Products Images

