







ASL Rabbit pAb

Catalog No YP-Ab-17873 Isotype IgG Reactivity Human,Mouse,Rat Applications IHC-P Gene Name ASL Alternative Names ASAL Research Field Signal Transduction Product Categories Primary antibody Host Rabbit Molecular Weight - Clonality Polyclonal Antibody Clonality No. - Dilution IHC: 1/50-1/100 Immunogen Fusion protein of human ASL Purification Affinity Purified Conjugation Unconjugated Modification Unmodified Form Liquid Buffer System pH 7.4 PBS, 0.05% NaN3, 40% Glycerol Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background This gene encodes a member of the lyase 1 family. The encoded protein forms a cytosolic homotetramer and primarily catalyzes the reversible hydrolytic cleavage of argininosuccinica into arginine and tumarate, an essential ste		
Reactivity Human,Mouse,Rat Applications IHC-P Gene Name ASL Alternative Names ASAL Research Field Signal Transduction Product Categories Primary antibody Host Rabbit Molecular Weight - Clonality Polyclonal Antibody Clonality No Dilution IHC: 1/50-1/100 Immunogen Fusion protein of human ASL Purification Affinity Purified Conjugation Unconjugated Modification Unmodified Form Liquid Buffer System pH 7.4 PBS, 0.05% NaN3, 40% Glycerol Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background Visene encodes a member of the lyase 1 family. The encoded protein forms a cytosolic homotetramer and primarily catalyzes the reversible hydrolytic cleavages and solicy as well as the visene and solicy as member of the lyase 1 family. The encoded protein forms a cytosolic homotetramer and primarily catalyzes the reversible hydrolytic cleavage autosomal recessive disorder argininosuccinia eind rumarate, an essential step in the liver in detoxifying ammonia via the urea cycle. Mutations in this gene result in the autosomal recessive disorder argininosuccinia caid runsarcine, and ensential step in the liver in detoxifying ammonia via the urea cycle. Mutations in this gene result in the autosomal recessive disorder argininosuccinia caid runsarcine, and ensential step in the liver in detoxifying ammonia via the urea cycle. Mutations in this gene result in the autosomal recessive disorder argininosuccinia caid runsarcine, acidence and efficiency. A nontranscribed pseudogene is also located on the long arm of chromosome 22. Alternatively spliced transcript variants encoding different	Catalog No	YP-Ab-17873
Applications IHC-P Gene Name ASL Alternative Names ASAL Research Field Signal Transduction Product Categories Primary antibody Host Rabbit Molecular Weight - Clonality Polyclonal Antibody Clonality No Dilution IHC: 1/50-1/100 Immunogen Fusion protein of human ASL Purification Affinity Purified Conjugation Unconjugated Modification Unmodified Form Liquid Buffer System pH 7.4 PBS, 0.05% NaN3, 40% Glycerol Concentration 1 mg/ml Purity ≥90% Storage Storage 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background This gene encodes a member of the lyase 1 family. The encoded protein forms a cytosolic homotetramer and primarily catalyzes the reversible hydrolytic cleavage of argininosuccinate into arginine and fumarate, an essential step in the liver in detoxifying ammonia via the urea cycle. Mutations in this gene result in the autosomal recessive disorder argininosuccinica caid lyase deficiency. A nontranscribed pseudogene is also located on the long arm of chromosome 22. Alternatively spilect transcript variants encoding different	Isotype	IgG
Gene Name ASL Alternative Names ASAL Research Field Signal Transduction Product Categories Primary antibody Host Rabbit Molecular Weight - Clonality Polyclonal Antibody Clonality No Dilution IHC: 1/50-1/100 Immunogen Fusion protein of human ASL Purification Affinity Purified Conjugation Unconjugated Modification Unmodified Form Liquid Buffer System pH 7.4 PBS, 0.05% NaN3, 40% Glycerol Concentration 1 mg/ml Purity ≥90% Storage Storage 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background This gene encodes a member of the lyase 1 family. The encoded protein forms a cytosolic homotetramer and primarily catalyzes the reversible hydrolytic cleavage of argininosuccinate into arginine and fumarate, an essential step in the liver in detoxifying ammonia via the urea cycle. Mutations in this gene result in the autosomal recessive disorder argininosuccinica acid lyase deficiency. A nontranscribed pseudogene is also located on the long arm of chromosome 22. Alternatively spilecet transcript variants encoding different	Reactivity	Human,Mouse,Rat
Alternative Names ASAL Research Field Signal Transduction Product Categories Primary antibody Host Rabbit Molecular Weight - Clonality Polyclonal Antibody Clonality No Dilution IHC: 1/50-1/100 Immunogen Fusion protein of human ASL Purification Affinity Purified Conjugation Unconjugated Modification Unmodified Form Liquid Buffer System pH 7.4 PBS, 0.05% NaN3, 40% Glycerol Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background This gene encodes a member of the Iyase 1 family. The encoded protein forms a cytosolic homotetramer and primarily catalyzes the reversible hydrolytic cleavage of argininosuccinate into arginine and fumarate, an essential step in the liver in deloxifying ammonia via the urea cycle. Mutations in this gene result in the autosomal recessive disorder argininosuccinic aciduria, or argininosuccinic acid lyase deficiency. A nontranscribed pseudogene is also located on the long arm of chromosome 22. Alternatively spliced transcript variants encoding different	Applications	IHC-P
Research Field Signal Transduction Product Categories Primary antibody Host Rabbit Molecular Weight - Clonality Polyclonal Antibody Clonality No. - Dilution IHC: 1/50-1/100 Immunogen Fusion protein of human ASL Purification Affinity Purified Conjugation Unconjugated Modification Unmodified Form Liquid Buffer System pH 7.4 PBS, 0.05% NaN3, 40% Glycerol Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background This gene encodes a member of the lyase 1 family. The encoded protein forms a cytosolic homotetramer and primarily catalyzes the reversible hydrolytic cleavage of argininosuccinate into arginine and fumartae, an essential step in the liver in detoxifying ammonia via the urea cycle. Mutations in this gene result in the autosomal recessive disorder argininosuccinate aciduria, or argininosuccinate acidoria, or argininosuccinic acid lyses deficiency. A nontranscribed pseudogene is also located on the long arm of chromosome 22. Alternatively spliced transcript variants encoding different	Gene Name	ASL
Product Categories Primary antibody Host Rabbit Molecular Weight - Clonality Polyclonal Antibody Clonality No Dilution IHC: 1/50-1/100 Immunogen Fusion protein of human ASL Purification Affinity Purified Conjugation Unconjugated Modification Unmodified Form Liquid Buffer System pH 7.4 PBS, 0.05% NaN3, 40% Glycerol Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background This gene encodes a member of the lyase 1 family. The encoded protein forms a cytosolic homotetramer and primarily catalyzes the reversible hydrolytic cleavage of argininosuccinate into arginine and furnarate, an essential step in the liver in detoxifying ammonia via the urea cycle. Mutations in this gene result in the autosomal recessive disorder argininosuccinic aciduly, ase deficiency. A nontranscribed pseudogene is also located on the long arm of chromosome 22. Alternatively spliced transcript variants encoding different	Alternative Names	ASAL
Host Rabbit Molecular Weight - Clonality Polyclonal Antibody Clonality No. - Dilution IHC: 1/50-1/100 Immunogen Fusion protein of human ASL Purification Affinity Purified Conjugation Unconjugated Modification Unmodified Form Liquid Buffer System pH 7.4 PBS, 0.05% NaN3, 40% Glycerol Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background This gene encodes a member of the lyase 1 family. The encoded protein forms a cytosolic homotetramer and primarily catalyzes the reversible hydrolytic cleavage of argininosuccinate into arginine and fumarate, an essential step in the liver in detoxifying ammonia via the urea cycle. Mutations in this gene result in the autosomal recessive disorder argininosuccinic acidity are deficiency. A nontranscribed pseudogene is also located on the long arm of chromosome 22. Alternatively spliced transcript variants encoding different	Research Field	Signal Transduction
Molecular Weight - Clonality Polyclonal Antibody Clonality No. - Dilution IHC: 1/50-1/100 Immunogen Fusion protein of human ASL Purification Affinity Purified Conjugation Unconjugated Modification Unmodified Form Liquid Buffer System pH 7.4 PBS, 0.05% NaN3, 40% Glycerol Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background This gene encodes a member of the Iyase 1 family. The encoded protein forms a cycles withutations in this gene result in the autosomal recessive disorder argininosuccinica cidultria, or argininosuccinic acid lypase deficiency. A nontranscribed pseudogene is also located on the long arm of chromosome 22. Alternatively spliced transcript variants encoding different	Product Categories	Primary antibody
Clonality No Dilution IHC: 1/50-1/100 Immunogen Fusion protein of human ASL Purification Affinity Purified Conjugation Unconjugated Modification Unmodified Form Liquid Buffer System pH 7.4 PBS, 0.05% NaN3, 40% Glycerol Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background This gene encodes a member of the lyase 1 family. The encoded protein forms a cytosolic homotetramer and primarily catalyzes the reversible hydrolytic cleavage of argininosuccinate into arginine and furnarate, an essential step in the liver in detoxifying ammonia via the urea cycle. Mutations in this gene result in the autosomal recessive disorder argininosuccinic acidiuria, or argininosuccinic a	Host	Rabbit
Clonality No. - Dilution IHC: 1/50-1/100 Immunogen Fusion protein of human ASL Purification Affinity Purified Conjugation Unconjugated Modification Unmodified Form Liquid Buffer System pH 7.4 PBS, 0.05% NaN3, 40% Glycerol Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background This gene encodes a member of the lyase 1 family. The encoded protein forms a cytosolic homotetramer and primarily catalyzes the reversible hydrolytic cleavage of argininosuccinale into arginine and fumarate, an essential step in the liver in detoxifying ammonia via the urea cycle. Mutations in this gene result in the autosomal recessive disorder argininosuccinic aciduria, or argininosuccinic acid lyase deficiency. A nontranscribed pseudogene is also located on the long arm of chromosome 22. Alternatively spiliced transcript variants encoding different	Molecular Weight	-
Dilution IHC: 1/50-1/100 Immunogen Fusion protein of human ASL Purification Affinity Purified Conjugation Unconjugated Modification Unmodified Form Liquid Buffer System pH 7.4 PBS, 0.05% NaN3, 40% Glycerol Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background This gene encodes a member of the lyase 1 family. The encoded protein forms a cytosolic homotetramer and primarily catalyzes the reversible hydrolytic cleavage of argininosuccinate into arginine and fumarate, an essential step in the liver in detoxifying ammonia via the urea cycle. Mutations in this gene result in the autosomal recessive disorder argininosuccinic aciduria, or argininosuccinic acid lyase deficiency. A nontranscribed pseudogene is also located on the long arm of chromosome 22. Alternatively spliced transcript variants encoding different	Clonality	Polyclonal Antibody
Immunogen Fusion protein of human ASL Purification Affinity Purified Conjugation Unconjugated Modification Unmodified Form Liquid Buffer System pH 7.4 PBS, 0.05% NaN3, 40% Glycerol Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background This gene encodes a member of the lyase 1 family. The encoded protein forms a cytosolic homotetramer and primarily catalyzes the reversible hydrolytic cleavage of argininosuccinate into arginine and fumarate, an essential step in the liver in detoxifying ammonia via the urea cycle. Mutations in this gene result in the autosomal recessive disorder argininosuccinic aciduria, or argininosuccinic acid lyase deficiency. A nontranscribed pseudogene is also located on the long arm of chromosome 22. Alternatively spliced transcript variants encoding different	Clonality No.	-
Purification Affinity Purified Conjugation Unconjugated Modification Unmodified Form Liquid Buffer System pH 7.4 PBS, 0.05% NaN3, 40% Glycerol Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background This gene encodes a member of the lyase 1 family. The encoded protein forms a cytosolic homotetramer and primarily catalyzes the reversible hydrolytic cleavage of argininosuccinate into arginine and furmarate, an essential step in the liver in detoxifying ammonia via the urea cycle. Mutations in this gene result in the autosomal recessive disorder argininosuccinic aciduria, or argininosuccinic acid lyase deficiency. A nontranscribed pseudogene is also located on the long arm of chromosome 22. Alternatively spliced transcript variants encoding different	Dilution	IHC: 1/50-1/100
Conjugation Unconjugated Modification Unmodified Form Liquid Buffer System pH 7.4 PBS, 0.05% NaN3, 40% Glycerol Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background This gene encodes a member of the lyase 1 family. The encoded protein forms a cytosolic homotetramer and primarily catalyzes the reversible hydrolytic cleavage of argininosuccinate into arginine and fumarate, an essential step in the liver in detoxifying ammonia via the urea cycle. Mutations in this gene result in the autosomal recessive disorder argininosuccinic aciduria, or argininosuccinic acid lyase deficiency. A nontranscribed pseudogene is also located on the long arm of chromosome 22. Alternatively spliced transcript variants encoding different	Immunogen	Fusion protein of human ASL
Modification Unmodified Form Liquid Buffer System pH 7.4 PBS, 0.05% NaN3, 40% Glycerol Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background This gene encodes a member of the lyase 1 family. The encoded protein forms a cytosolic homotetramer and primarily catalyzes the reversible hydrolytic cleavage of argininosuccinate into arginine and fumarate, an essential step in the liver in detoxifying ammonia via the urea cycle. Mutations in this gene result in the autosomal recessive disorder argininosuccinic aciduria, or argininosuccinic acid lyase deficiency. A nontranscribed pseudogene is also located on the long arm of chromosome 22. Alternatively spliced transcript variants encoding different	Purification	Affinity Purified
Form Liquid Buffer System pH 7.4 PBS, 0.05% NaN3, 40% Glycerol Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background This gene encodes a member of the lyase 1 family. The encoded protein forms a cytosolic homotetramer and primarily catalyzes the reversible hydrolytic cleavage of argininosuccinate into arginine and fumarate, an essential step in the liver in detoxifying ammonia via the urea cycle. Mutations in this gene result in the autosomal recessive disorder argininosuccinic aciduria, or argininosuccinic acid lyase deficiency. A nontranscribed pseudogene is also located on the long arm of chromosome 22. Alternatively spliced transcript variants encoding different	Conjugation	Unconjugated
Buffer System Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background This gene encodes a member of the lyase 1 family. The encoded protein forms a cytosolic homotetramer and primarily catalyzes the reversible hydrolytic cleavage of argininosuccinate into arginine and fumarate, an essential step in the liver in detoxifying ammonia via the urea cycle. Mutations in this gene result in the autosomal recessive disorder argininosuccinic aciduria, or argininosuccinic acid lyase deficiency. A nontranscribed pseudogene is also located on the long arm of chromosome 22. Alternatively spliced transcript variants encoding different	Modification	Unmodified
Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background This gene encodes a member of the lyase 1 family. The encoded protein forms a cytosolic homotetramer and primarily catalyzes the reversible hydrolytic cleavage of argininosuccinate into arginine and fumarate, an essential step in the liver in detoxifying ammonia via the urea cycle. Mutations in this gene result in the autosomal recessive disorder argininosuccinic aciduria, or argininosuccinic acid lyase deficiency. A nontranscribed pseudogene is also located on the long arm of chromosome 22. Alternatively spliced transcript variants encoding different	Form	Liquid
Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background This gene encodes a member of the lyase 1 family. The encoded protein forms a cytosolic homotetramer and primarily catalyzes the reversible hydrolytic cleavage of argininosuccinate into arginine and fumarate, an essential step in the liver in detoxifying ammonia via the urea cycle. Mutations in this gene result in the autosomal recessive disorder argininosuccinic aciduria, or argininosuccinic acid lyase deficiency. A nontranscribed pseudogene is also located on the long arm of chromosome 22. Alternatively spliced transcript variants encoding different	Buffer System	pH 7.4 PBS, 0.05% NaN3, 40% Glycerol
Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background This gene encodes a member of the lyase 1 family. The encoded protein forms a cytosolic homotetramer and primarily catalyzes the reversible hydrolytic cleavage of argininosuccinate into arginine and fumarate, an essential step in the liver in detoxifying ammonia via the urea cycle. Mutations in this gene result in the autosomal recessive disorder argininosuccinic aciduria, or argininosuccinic acid lyase deficiency. A nontranscribed pseudogene is also located on the long arm of chromosome 22. Alternatively spliced transcript variants encoding different	Concentration	1 mg/ml
Cycles. Background This gene encodes a member of the lyase 1 family. The encoded protein forms a cytosolic homotetramer and primarily catalyzes the reversible hydrolytic cleavage of argininosuccinate into arginine and fumarate, an essential step in the liver in detoxifying ammonia via the urea cycle. Mutations in this gene result in the autosomal recessive disorder argininosuccinic aciduria, or argininosuccinic acid lyase deficiency. A nontranscribed pseudogene is also located on the long arm of chromosome 22. Alternatively spliced transcript variants encoding different	Purity	≥90%
cytosolic homotetramer and primarily catalyzes the reversible hydrolytic cleavage of argininosuccinate into arginine and fumarate, an essential step in the liver in detoxifying ammonia via the urea cycle. Mutations in this gene result in the autosomal recessive disorder argininosuccinic aciduria, or argininosuccinic acid lyase deficiency. A nontranscribed pseudogene is also located on the long arm of chromosome 22. Alternatively spliced transcript variants encoding different	Storage	·
	Background	cytosolic homotetramer and primarily catalyzes the reversible hydrolytic cleavage of argininosuccinate into arginine and fumarate, an essential step in the liver in detoxifying ammonia via the urea cycle. Mutations in this gene result in the autosomal recessive disorder argininosuccinic aciduria, or argininosuccinic acid lyase deficiency. A nontranscribed pseudogene is also located on the long arm of chromosome 22. Alternatively spliced transcript variants encoding different



UpingBio technology Co.,Ltd

€ Tel: 400-999-8863 🖷 Email:UpingBio@163.com



matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

