





MGAT2 rabbit pAb

| 货号 | YP-Ab-17965 |
|--------|---|
| 同位型 | IgG |
| 应用 | WB |
| 种属 | Human;Mouse;Rat |
| 靶点 | MGAT2 |
| 基因名称 | MGAT2 |
| 蛋白名称 | Alpha-1,6-mannosyl-glycoprotein 2-beta-N-acetylglucosaminyltransferase (EC 2.4.1.143) (Beta-1,2-N-acetylglucosaminyltransferase II) (GlcNAc-T II) (GNT-II) (Mannoside acetylglucosaminyltransferase 2) |
| 免疫原 | Synthesized peptide derived from human MGAT2 |
| 特异性 | This antibody detects endogenous levels of MGAT2 at Human, Mouse,Rat |
| 组成 | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| 来源 | Polyclonal, Rabbit,IgG |
| 稀释 | WB 1:500-2000 |
| 纯化工艺 | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| 分子量 | 49kD |
| 功能 | Plays an essential role in protein N-glycosylation. Catalyzes the transfer of N-acetylglucosamine (GlcNAc) onto the free terminal mannose moiety in the core structure of the nascent N-linked glycan chain, giving rise to the second branch in complex glycans. |
| 细胞定位 | Golgi apparatus membrane ; Single-pass type II membrane protein . |
| 浓度 | 1 mg/ml |
| 储存 | -15°C to -25°C/1 year(Do not lower than -25°C) |
| 有关注意事项 | Avoid repeated freezing and thawing! |
| 使用建议 | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. |
| | |

Products Images