



TNIP1 rabbit pAb

货号	YP-Ab-17972
同位型	IgG
应用	WB
种属	Human;Mouse
靶点	TNIP1
基因名称	TNIP1 KIAA0113 NAF1
蛋白名称	TNFAIP3-interacting protein 1 (A20-binding inhibitor of NF-kappa-B activation 1) (ABIN-1) (HIV-1 Nef-interacting protein) (Nef-associated factor 1) (Naf1) (Nip40-1) (Virion-associated nuclear shuttlin
免疫原	Synthesized peptide derived from human TNIP1
特异性	This antibody detects endogenous levels of TNIP1 at Human, Mouse
组成	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
来源	Polyclonal, Rabbit,IgG
稀释	WB 1:500-2000
纯化工艺	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
分子量	70kD
功能	Inhibits NF-kappa-B activation and TNF-induced NF-kappa-B-dependent gene expression by regulating A20/TNFAIP3-mediated deubiquitination of IKBKG; proposed to link A20/TNFAIP3 to ubiquitinated IKBKG. Involved in regulation of EGF-induced ERK1/ERK2 signaling pathway; blocks MAPK3/MAPK1 nuclear translocation and MAPK1-dependent transcription. Increases cell surface CD4(T4) antigen expression. Involved in the anti-inflammatory response of macrophages and positively regulates TLR-induced activation of CEBPB. Involved in the prevention of autoimmunity; this function implicates binding to polyubiquitin. Involved in leukocyte integrin activation during inflammation; this function is mediated by association with SELPLG and dependent on phosphorylation by SRC-family kinases. Interacts with HIV-1 matrix protein and is packaged into virions and overexpression can inhibit viral replication. May regu
细胞定位	Cytoplasm. Nucleus. Shuttles between the nucleus and cytoplasm in a CRM1-dependent manner.
组织表达	Ubiquitous. Strongly expressed in peripheral blood lymphocytes, spleen and skeletal muscle, and is weakly expressed in the brain. In peripheral blood mononucleocytes, isoform 4 is mainly expressed and isoform 1 and isoform 7 are almost not expressed. Expression of isoform 1 and isoform 7 increases in leukemic cells.
浓度	1 mg/ml
储存	-15°C to -25°C/1 year(Do not lower than -25°C)



有关注意事项

Avoid repeated freezing and thawing!

使用建议

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images