

## MED19 rabbit pAb

|        |  |
|--------|--|
| 货号     | YP-Ab-18018  |
| 同位型    | IgG  |
| 应用     | WB   |
| 种属     | Human;Mouse  |
| 靶点     | MED19  |
| 基因名称   | MED19 LCMR1  |
| 蛋白名称   | Mediator of RNA polymerase II transcription subunit 19 (Lung cancer metastasis-related protein 1) (Mediator complex subunit 19)  |
| 免疫原    | Synthesized peptide derived from human MED19   |
| 特异性    | This antibody detects endogenous levels of MED19 at Human, Mouse   |
| 组成     | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |
| 来源     | Polyclonal, Rabbit,IgG   |
| 稀释     | WB 1:500-2000  |
| 纯化工艺   | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  |
| 分子量    | 27kD   |
| 功能     | Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors. |
| 细胞定位   | Nucleus .  |
| 浓度     | 1 mg/ml  |
| 储存     | -15°C to -25°C/1 year(Do not lower than -25°C)   |
| 有关注意事项 | Avoid repeated freezing and thawing!   |
| 使用建议   | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.  |



## Products Images