

Website: www.upingBio.com

ARHGEF18 rabbit pAb

货号	YP-Ab-18050
同位型	lgG
应用	WB
种属	Human;Mouse
靶点	ARHGEF18
基因名称	ARHGEF18 KIAA0521
蛋白名称	Rho guanine nucleotide exchange factor 18 (114 kDa Rho-specific guanine nucleotide exchange factor) (p114-Rho-GEF) (p114RhoGEF) (Septin-associated RhoGEF) (SA-RhoGEF)
免疫原	Synthesized peptide derived from human ARHGEF18
特异性	This antibody detects endogenous levels of ARHGEF18 at Human, Mouse
组成	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
来源	Polyclonal, Rabbit,IgG
稀释	WB 1:500-2000
纯化工艺	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
分子量	150kD
功能	Acts as guanine nucleotide exchange factor (GEF) for RhoA GTPases. Its activation induces formation of actin stress fibers. Also acts as a GEF for RAC1, inducing production of reactive oxygen species (ROS). Does not act as a GEF for CDC42. The G protein beta-gamma (Gbetagamma) subunits of heterotrimeric G proteins act as activators, explaining the integrated effects of LPA and other G-protein coupled receptor agonists on actin stress fiber formation, cell shape change and ROS production. Required for EPB41L4B-mediated regulation of the circumferential actomyosin belt in epithelial cells.
细胞定位	Cytoplasm . Cytoplasm, cytoskeleton . Cell membrane . Apical cell membrane . In unactivated eosinophils, distributed around the cell periphery in the perimembranous region (PubMed:29601110). In activated eosinophils, relocates to the tip of the nucleopod, a membrane structure formed during activation when the nucleus moves to one end of the cell, and is also concentrated in membrane protrusions at the opposite end of the cell (PubMed:29601110). Localizes to the apical cell membrane in epithelial cells (PubMed:22006950).
组织表达	Expressed in all tissues tested with highest expression in kidney and pancreas. Weakly or not expressed in liver, skeletal muscle and testis. Isoform 1: Expressed in eosinophils (PubMed:29601110). Isoform 2: Expressed in eosinophils (PubMed:29601110). Isoform 3: Expressed in eosinophils (PubMed:29601110). Isoform 4: Not detected in eosinophils (PubMed:29601110).
浓度	1 mg/ml
储存	-15°C to -25°C/1 year(Do not lower than -25°C)



🔇 Tel: 400-999-8863 💌 Email:UpingBio@163.com

@ Website: www.upingBio.com

有关注意事项

Avoid repeated freezing and thawing!

使用建议

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images