



## PARVB rabbit pAb

| 货号     | YP-Ab-18060  |
|--------|--|
| 同位型    | IgG  |
| 应用     | WB   |
| 种属     | Human;Mouse  |
| 靶点     | PARVB  |
| 基因名称   | PARVB CGI-56   |
| 蛋白名称   | Beta-parvin (Affixin)  |
| 免疫原    | Synthesized peptide derived from human PARVB   |
| 特异性    | This antibody detects endogenous levels of PARVB at Human, Mouse   |
| 组成     | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |
| 来源     | Polyclonal, Rabbit,IgG   |
| 稀释     | WB 1:500-2000  |
| 纯化工艺   | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  |
| 分子量    | 40kD   |
| 功能     | Adapter protein that plays a role in integrin signaling via ILK and in activation of<br>the GTPases CDC42 and RAC1 by guanine exchange factors, such as<br>ARHGEF6. Is involved in the reorganization of the actin cytoskeleton and<br>formation of lamellipodia. Plays a role in cell adhesion, cell spreading,<br>establishment or maintenance of cell polarity, and cell migration. |
| 细胞定位   | Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein;<br>Cytoplasmic side. Cytoplasm, cytoskeleton. Cell projection, lamellipodium.<br>Cytoplasm, myofibril, sarcomere. Cytoplasm, myofibril, sarcomere, Z line.<br>Constituent of focal adhesions. Detected at the tips of the leading edge of cells.<br>Colocalizes with F-actin at the tips of lamellipodia.   |
| 组织表达   | Expressed predominantly in heart and skeletal muscle.  |
| 浓度     | 1 mg/ml  |
| 储存     | -15°C to -25°C/1 year(Do not lower than -25°C)   |
| 有关注意事项 | Avoid repeated freezing and thawing!   |
| 使用建议   | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.  |
|        |  |



🕔 Tel: 400-999-8863 💌 Email:UpingBio@163.com



**Products Images**