

LPCAT1 rabbit pAb

Catalog No	YP-Ab-18072
Isotype	IgG
Reactivity	WB
Applications	Human;Mouse;Rat
Gene Name	LPCAT1 AYTL2 PFAAP3
Target	LPCAT1
Research Field	Lysophosphatidylcholine acyltransferase 1 (LPC acyltransferase 1) (LPCAT-1) (LysoPC acyltransferase 1) (EC 2.3.1) (1-acylglycerophosphocholine O-acyltransferase) (EC 2.3.1.23) (1-alkylglycerophospho
Specificity	This antibody detects endogenous levels of LPCAT1 at Human, Mouse,Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Molecular Weight	59kD
Source	Polyclonal, Rabbit,IgG
Dilution	WB 1:500-2000
Immunogen	Synthesized peptide derived from human LPCAT1
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Function	Exhibits acyltransferase activity . Exhibits acetyltransferase activity (By similarity). Activity is calcium-independent (By similarity). Catalyzes the conversion of lysophosphatidylcholine (1-acyl-sn-glycero-3-phosphocholine or LPC) into phosphatidylcholine (1,2-diacyl-sn-glycero-3-phosphocholine or PC) . Catalyzes the conversion 1-acyl-sn-glycerol-3-phosphate (lysophosphatidic acid or LPA) into 1,2-diacyl-sn-glycerol-3-phosphate (lysophosphatidic acid or LPA) into 1,2-diacyl-sn-glycerol-3-phosphate (phosphatidic acid or PA) by incorporating an acyl moiety at the sn-2 position of the glycerol backbone (By similarity). Displays a clear preference for saturated fatty acyl-CoAs, and 1-myristoyl or 1-palmitoyl LPC as acyl donors and acceptors, respectively (By similarity). Involved in platelet-activating factor (PAF) biosynthesis by catalyzing the conversion of the PAF precursor, 1-O-alkyl-sn-glycero-3-phosphocholine (lyso-PAF) into 1-O-alkyl-2-acetyl-sn-glycero-3-phosphocholine (lyso-PAF) into
Subcellular Location	Endoplasmic reticulum membrane ; Single-pass type II membrane protein . Golgi apparatus membrane ; Single-pass type II membrane protein . Cell membrane ; Single-pass type II membrane
Concentration	1 mg/ml
Purity	≥90%
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Background	



Avoid repeated freezing and thawing!

🕚 Tel: 400-999-8863 💌 Email:UpingBio@163.com



matters needing attention

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images