



ORAI1 rabbit pAb

货号	YP-Ab-18121
同位型	IgG
应用	WB
种属	Human;Mouse;Rat
靶点	ORAI1
基因名称	ORAI1 CRACM1 TMEM142A
蛋白名称	Calcium release-activated calcium channel protein 1 (Protein orai-1) (Transmembrane protein 142A)
免疫原	Synthesized peptide derived from human ORAI1
特异性	This antibody detects endogenous levels of ORAI1 at Human, Mouse,Rat
组成	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
来源	Polyclonal, Rabbit,IgG
稀释	WB 1:500-2000
纯化工艺	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
分子量	33kD
功能	Ca(2+) release-activated Ca(2+) (CRAC) channel subunit which mediates Ca(2+) influx following depletion of intracellular Ca(2+) stores and channel activation by the Ca(2+) sensor, STIM1 . CRAC channels are the main pathway for Ca(2+) influx in T-cells and promote the immune response to pathogens by activating the transcription factor NFAT . Plays a prominent role in Ca(2+) influx at the basolateral membrane of mammary epithelial cells independently of the Ca(2+) content of endoplasmic reticulum or Golgi stores. May mediate transepithelial transport of large quantities of Ca(2+) for milk secretion.
细胞定位	Cell membrane ; Multi-pass membrane protein . Basolateral cell membrane ; Multi-pass membrane protein . Isoform beta is more mobile in the plasma membrane (PubMed:23307288). Colocalizes with STIM1 at the cell membrane (PubMed:27185316). .
组织表达	Expressed in naive CD4 and CD8 T cells (at protein level) (PubMed:26956484). Expressed at similar levels in naive and effector T helper cells (PubMed:20354224).
浓度	1 mg/ml
储存	-15°C to -25°C/1 year(Do not lower than -25°C)
有关注意事项	Avoid repeated freezing and thawing!
使用建议	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images