

🔇 Tel: 400-999-8863 💌 Email:UpingBio@163.com



PDI rabbit pAb

货号	YP-Ab-18190
同位型	IgG
应用	WB
种属	Human;Mouse;Rat
靶点	P4HB
基因名称	P4HB ERBA2L PDI PDIA1 PO4DB
蛋白名称	Protein disulfide-isomerase (PDI) (EC 5.3.4.1) (Cellular thyroid hormone-binding protein) (Prolyl 4-hydroxylase subunit beta) (p55)
免疫原	Synthesized peptide derived from human PDI
特异性	This antibody detects endogenous levels of PDI at Human, Mouse,Rat
组成	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
来源	Polyclonal, Rabbit,IgG
稀释	WB 1:500-2000
纯化工艺	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
分子量	56kD
功能	This multifunctional protein catalyzes the formation, breakage and rearrangement of disulfide bonds. At the cell surface, seems to act as a reductase that cleaves disulfide bonds of proteins attached to the cell. May therefore cause structural modifications of exofacial proteins. Inside the cell, seems to form/rearrange disulfide bonds of nascent proteins. At high concentrations, functions as a chaperone that inhibits aggregation of misfolded proteins. At low concentrations, facilitates aggregation (anti-chaperone activity). May be involved with other chaperones in the structural modification of the TG precursor in hormone biogenesis. Also acts a structural subunit of various enzymes such as prolyl 4-hydroxylase and microsomal triacylglycerol transfer protein MTTP. Receptor for LGALS9; the interaction retains P4HB at the cell surface of Th2 T helper cells, increasing disulfide reductase
细胞定位	Endoplasmic reticulum . Endoplasmic reticulum lumen . Melanosome . Cell membrane ; Peripheral membrane protein . Highly abundant. In some cell types, seems to be also secreted or associated with the plasma membrane, where it undergoes constant shedding and replacement from intracellular sources (Probable). Localizes near CD4-enriched regions on lymphoid cell surfaces (PubMed
浓度	1 mg/ml
储存	-15°C to -25°C/1 year(Do not lower than -25°C)
有关注意事项	Avoid repeated freezing and thawing!



UpingBio technology Co.,Ltd

C Tel: 400-999-8863 💌 Email:UpingBio@163.com

Website: www.upingBio.com

使用建议

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images