



PIEZ1 rabbit pAb

货号	YP-Ab-18283
同位型	IgG
应用	IHC;WB
种属	Human;Mouse;Rat
靶点	PIEZ1
基因名称	PIEZO1 FAM38A KIAA0233
蛋白名称	Piezo-type mechanosensitive ion channel component 1 (Membrane protein induced by beta-amyloid treatment) (Mib) (Protein FAM38A)
免疫原	Synthesized peptide derived from human C-terminal PIEZ1
特异性	This antibody detects endogenous levels of PIEZ1 at Human, Mouse,Rat
组成	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
来源	Polyclonal, Rabbit,IgG
稀释	WB 1:500-2000 IHC 1:50-200
纯化工艺	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
其他名称	Piezo-type mechanosensitive ion channel component 1 (Membrane protein induced by beta-amyloid treatment) (Mib) (Protein FAM38A)
分子量	277kD
功能	Pore-forming subunit of a mechanosensitive non-specific cation channel . Generates currents characterized by a linear current-voltage relationship that are sensitive to ruthenium red and gadolinium. Plays a key role in epithelial cell adhesion by maintaining integrin activation through R-Ras recruitment to the ER, most probably in its activated state, and subsequent stimulation of calpain signaling . In the kidney, may contribute to the detection of intraluminal pressure changes and to urine flow sensing. Acts as shear-stress sensor that promotes endothelial cell organization and alignment in the direction of blood flow through calpain activation . Plays a key role in blood vessel formation and vascular structure in both development and adult physiology (By similarity). Acts as sensor of phosphatidylserine (PS) flipping at the plasma membrane and governs morphogenesis of muscle cells. I
细胞定位	Endoplasmic reticulum membrane ; Multi-pass membrane protein. Endoplasmic reticulum-Golgi intermediate compartment membrane . Cell membrane ; Multi-pass membrane protein . Cell projection, lamellipodium membrane . Cell membrane ; Multi-pass membrane protein. In erythrocytes, located in the plasma membrane (PubMed:22529292, PubMed:23479567). Accumulates at the leading apical lamellipodia of endothelial cells in response to shear stress (PubMed:25119035). Colocalizes with F-actin and MYH9 at the actomyosin cortex in myoblasts. .
组织表达	Expressed in numerous tissues. In normal brain, expressed exclusively in neurons, not in astrocytes. In Alzheimer disease brains, expressed in about half of the activated astrocytes located around classical senile plaques. In Parkinson disease substantia nigra, not detected in melanin-containing neurons nor in

activated astrocytes. Expressed in erythrocytes (at protein level). Expressed in myoblasts (at protein level).

浓度 1 mg/ml

储存 -15°C to -25°C/1 year(Do not lower than -25°C)

有关注意事项 Avoid repeated freezing and thawing!

使用建议 This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images