







ESCO2 rabbit pAb

| 货号 | YP-Ab-18295 |
|--------------------------------|---|
| 同位型 | IgG |
| 应用 | WB |
| 种属 | Human;Mouse |
| 靶点 | ESCO2 |
| 基因名称 | ESCO2 |
| 蛋白名称 | N-acetyltransferase ESCO2 (EC 2.3.1) (Establishment of cohesion 1 homolog 2) (ECO1 homolog 2) |
| 免疫原 | Synthesized peptide derived from human ESCO2 |
| 特异性 | This antibody detects endogenous levels of ESCO2 at Human, Mouse |
| 组成 | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| 来源 | Polyclonal, Rabbit,IgG |
| 稀释 | WB 1:500-2000 |
| 纯化工艺 | The antibody was affinity-purified from rabbit antiserum by |
| | affinity-chromatography using epitope-specific immunogen. |
| 分子量 | affinity-chromatography using epitope-specific immunogen. 66kD |
| 分子量功能 | |
| | Acetyltransferase required for the establishment of sister chromatid cohesion. Couples the processes of cohesion and DNA replication to ensure that only sister chromatids become paired together. In contrast to the structural cohesins, the deposition and establishment factors are required only during the S phase. |
| 功能 | Acetyltransferase required for the establishment of sister chromatid cohesion . Couples the processes of cohesion and DNA replication to ensure that only sister chromatids become paired together. In contrast to the structural cohesins, the deposition and establishment factors are required only during the S phase. Acetylates the cohesin component SMC3 . Nucleus . Chromosome . Nuclear in interphase cells, excluded from chromosomes during metaphase but reassociates with chromosomes in |
| 功能 细胞定位 | Acetyltransferase required for the establishment of sister chromatid cohesion. Couples the processes of cohesion and DNA replication to ensure that only sister chromatids become paired together. In contrast to the structural cohesins, the deposition and establishment factors are required only during the S phase. Acetylates the cohesin component SMC3. Nucleus. Chromosome. Nuclear in interphase cells, excluded from chromosomes during metaphase but reassociates with chromosomes in telophase. Widely expressed in fetal tissues. In adult, it is expressed in thymus, placenta and |
| 功能 细胞定位 组织表达 | Acetyltransferase required for the establishment of sister chromatid cohesion. Couples the processes of cohesion and DNA replication to ensure that only sister chromatids become paired together. In contrast to the structural cohesins, the deposition and establishment factors are required only during the S phase. Acetylates the cohesin component SMC3. Nucleus. Chromosome. Nuclear in interphase cells, excluded from chromosomes during metaphase but reassociates with chromosomes in telophase. Widely expressed in fetal tissues. In adult, it is expressed in thymus, placenta and small intestine. |
| 功能 细胞定位 组织表达 浓度 | Acetyltransferase required for the establishment of sister chromatid cohesion . Couples the processes of cohesion and DNA replication to ensure that only sister chromatids become paired together. In contrast to the structural cohesins, the deposition and establishment factors are required only during the S phase. Acetylates the cohesin component SMC3 . Nucleus . Chromosome . Nuclear in interphase cells, excluded from chromosomes during metaphase but reassociates with chromosomes in telophase. Widely expressed in fetal tissues. In adult, it is expressed in thymus, placenta and small intestine. |
| 功能 细胞定位 组织表达 浓度 储存 | Acetyltransferase required for the establishment of sister chromatid cohesion . Couples the processes of cohesion and DNA replication to ensure that only sister chromatids become paired together. In contrast to the structural cohesins, the deposition and establishment factors are required only during the S phase. Acetylates the cohesin component SMC3 . Nucleus . Chromosome . Nuclear in interphase cells, excluded from chromosomes during metaphase but reassociates with chromosomes in telophase Widely expressed in fetal tissues. In adult, it is expressed in thymus, placenta and small intestine. 1 mg/ml -15°C to -25°C/1 year(Do not lower than -25°C) |



UpingBio technology Co.,Ltd

€ Tel: 400-999-8863 🛎 Email:UpingBio@163.com



| Products Images | | |
|-----------------|--|--|
| | | |
| | | |
| | | |