

Ø Website: www.upingBio.com

## AIFL Polyclonal Antibody

| Catalog No         | YP-Ab-00295   |
|--------------------|---|
| lsotype            | lgG   |
| Reactivity         | Human;Mouse   |
| Applications       | WB;IHC;IF;ELISA   |
| Gene Name          | AIFM3   |
| Protein Name       | Apoptosis-inducing factor 3   |
| Immunogen          | The antiserum was produced against synthesized peptide derived from human AIFM3. AA range:10-59   |
| Specificity        | AIFL Polyclonal Antibody detects endogenous levels of AIFL protein.   |
| Formulation        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| Source             | Polyclonal, Rabbit,IgG  |
| Purification       | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.   |
| Dilution           | Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.<br>Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.   |
| Concentration      | 1 mg/ml   |
| Purity             | ≥90%  |
| Storage Stability  | -20°C/1 year  |
| Synonyms           | AIFM3; AIFL; Apoptosis-inducing factor 3; Apoptosis-inducing factor-like protein  |
| Observed Band      | 67kD  |
| Cell Pathway       | Mitochondrion . Does not translocate to the nucleus upon induction of apoptosis.  |
| Tissue Specificity | Ubiquitous. Expressed in bone marrow, cerebral cortex, liver, ovary, thymus, thyroid gland and tongue (at protein level).   |
| Function           | domain:The Rieske domain induces apoptosis.,function:Induces apoptosis<br>through a caspase dependent pathway. Reduces mitochondrial membrane<br>potential.,similarity:Belongs to the FAD-dependent oxidoreductase<br>family.,similarity:Contains 1 Rieske domain.,subcellular location:Does not<br>translocate to the nucleus upon induction of apoptosis.,tissue<br>specificity:Ubiquitous. Expressed in bone marrow, cerebral cortex, liver, ovary,<br>thymus, thyroid gland and tongue (at protein level)., |
| Background         | domain:The Rieske domain induces apoptosis.,function:Induces apoptosis<br>through a caspase dependent pathway. Reduces mitochondrial membrane<br>potential.,similarity:Belongs to the FAD-dependent oxidoreductase<br>family.,similarity:Contains 1 Rieske domain.,subcellular location:Does not<br>translocate to the nucleus upon induction of apoptosis.,tissue<br>specificity:Ubiquitous. Expressed in bone marrow, cerebral cortex, liver, ovary,  |



## UpingBio technology Co.,Ltd

🔇 Tel: 400-999-8863 📼 Emall:Upingbio.163.com

Website: www.upingBio.com

ingBio.com

## thymus, thyroid gland and tongue (at protein level).,

| matters needing attention | Avoid repeated freezing and thawing!  |
|---------------------------|---|
| Usage suggestions         | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. |

