




# DAP-1 Polyclonal Antibody

<b>Catalog No</b>	YP-Ab-00369
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	IHC;IF;ELISA
<b>Gene Name</b>	DAP
<b>Protein Name</b>	Death-associated protein 1
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human DAP. AA range:11-60
<b>Specificity</b>	DAP-1 Polyclonal Antibody detects endogenous levels of DAP-1 protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	IHC: 1/100 - 1/300. ELISA: 1/40000.. IF 1:50-200
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	DAP; DAP1; Death-associated protein 1; DAP-1
<b>Observed Band</b>	
<b>Cell Pathway</b>	
<b>Tissue Specificity</b>	Epithelium,Uterus,
<b>Function</b>	function:Involved in mediating interferon-gamma-induced cell death.,
<b>Background</b>	This gene encodes a basic, proline-rich, 15-kD protein. The protein acts as a positive mediator of programmed cell death that is induced by interferon-gamma. Alternatively spliced transcript variants encoding distinct isoforms have been found for this gene. [provided by RefSeq, May 2014],
<b>matters needing attention</b>	Avoid repeated freezing and thawing!
<b>Usage suggestions</b>	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue, using DAP Antibody. The picture on the right is blocked with the synthesized peptide.