

Tel: 400-999-8863
■ Emall:Upingbio.163.com





PSA Monoclonal Antibody

| Catalog No | YP-Ab-03416 |
|--------------------|--|
| Isotype | IgG |
| Reactivity | Human |
| Applications | WB;IHC;IF;FCM;ELISA |
| Gene Name | KLK3 |
| Protein Name | Prostate-specific antigen |
| Immunogen | Purified recombinant fragment of PSA (aa26-251) expressed in E. Coli. |
| Specificity | PSA Monoclonal Antibody detects endogenous levels of PSA protein. |
| Formulation | Ascitic fluid containing 0.03% sodium azide,0.5% BSA, 50%glycerol. |
| Source | Monoclonal, Mouse |
| Purification | Affinity purification |
| Dilution | WB: 1/500 - 1/2000. IHC: 1/200 - 1/1000. Flow cytometry: 1/200 - 1/400. ELISA: 1/10000 IF 1:50-200 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | KLK3; APS; Prostate-specific antigen; PSA; Gamma-seminoprotein; Seminin; Kallikrein-3; P-30 antigen; Semenogelase |
| Observed Band | |
| Cell Pathway | Secreted. |
| Tissue Specificity | Leukocyte,PNS,Prostate, |
| Function | catalytic activity:Preferential cleavage: -Tyr- -Xaa,function:Hydrolyzes semenogelin-1 thus leading to the liquefaction of the seminal coagulum.,online information:Prostate-specific antigen entry,similarity:Belongs to the peptidase S1 family.,similarity:Belongs to the peptidase S1 family. Kallikrein subfamily.,similarity:Contains 1 peptidase S1 domain., |
| Background | Kallikreins are a subgroup of serine proteases having diverse physiological functions. Growing evidence suggests that many kallikreins are implicated in carcinogenesis and some have potential as novel cancer and other disease biomarkers. This gene is one of the fifteen kallikrein subfamily members located in a cluster on chromosome 19. Its protein product is a protease present in seminal plasma. It is thought to function normally in the liquefaction of seminal coagulum, presumably by hydrolysis of the high molecular mass seminal vesicle protein. Serum level of this protein, called PSA in the clinical setting, is useful in the diagnosis and monitoring of prostatic carcinoma. Alternate splicing of this gene generates several transcript variants encoding different isoforms. [provided by |



UpingBio technology Co.,Ltd

C Tel: 400-999-8863 🗷 Email:Upingbio.163.com



RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


Products Images



Western Blot analysis using PSA Monoclonal Antibody against truncated PSA recombinant protein.



Immunohistochemistry analysis of paraffin-embedded human prostate carcinoma tissues, showing cytoplasmic localization with DAB staining using PSA Monoclonal Antibody.



Flow cytometric analysis of PC-3 cells using PSA Monoclonal Antibody (right) and negative control (left).