



HXK I Monoclonal Antibody

Catalog No	YP-Ab-14157
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;IHC;IF;FCM;ELISA
Gene Name	HK1
Protein Name	Hexokinase-1
Immunogen	Purified recombinant fragment of human HXK I expressed in E. Coli.
Specificity	HXK I Monoclonal Antibody detects endogenous levels of HXK I protein.
Formulation	Ascitic fluid containing 0.03% sodium azide,0.5% BSA, 50%glycerol.
Source	Monoclonal, Mouse
Purification	Affinity purification
Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/200 - 1/1000. Immunofluorescence: 1/200 - 1/1000. Flow cytometry: 1/200 - 1/400. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	HK1; Hexokinase-1; Brain form hexokinase; Hexokinase type I; HK I
Observed Band	
Cell Pathway	Mitochondrion outer membrane ; Peripheral membrane protein . Cytoplasm, cytosol . The mitochondrial-binding peptide (MBP) region promotes association with the mitochondrial outer membrane (Probable). Dissociates from the mitochondrial outer membrane following inhibition by N-acetyl-D-glucosamine, leading to relocation to the cytosol (PubMed:27374331). .
Tissue Specificity	Isoform 2: Erythrocyte specific (Ref.6). Isoform 3: Testis-specific (PubMed:10978502). Isoform 4: Testis-specific (PubMed:10978502).
Function	catalytic activity:ATP + D-hexose = ADP + D-hexose 6-phosphate.,disease:Defects in HK1 are the cause of hexokinase deficiency [MIM:235700]. Hexokinase deficiency is a rare autosomal recessive disease with nonspherocytic hemolytic anemia as the predominant clinical feature.,domain:The N- and C-terminal halves of this hexokinase show extensive sequence similarity to each other. The catalytic activity is associated with the C-terminus while regulatory function is associated with the N-terminus.,enzyme regulation:Hexokinase is an allosteric enzyme inhibited by its product Glc-6-P.,miscellaneous:In vertebrates there are four major glucose-phosphorylating isoenzymes, designated hexokinase I, II, III and IV (glucokinase).,online information:Hexokinase entry,pathway:Carbohydrate metabolism; hexose metabolism.,similarity:Belongs to the hexokinase



family.,subcellular location:Its hydrophobic N-ter

Background

Hexokinases phosphorylate glucose to produce glucose-6-phosphate, the first step in most glucose metabolism pathways. This gene encodes a ubiquitous form of hexokinase which localizes to the outer membrane of mitochondria. Mutations in this gene have been associated with hemolytic anemia due to hexokinase deficiency. Alternative splicing of this gene results in several transcript variants which encode different isoforms, some of which are tissue-specific. [provided by RefSeq, Apr 2016],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


Products Images




Western Blot analysis using HXK I Monoclonal Antibody against Jurkat (1), HeLa (2), HepG2 (3) and NIH/3T3 (4) cell lysate.



Immunohistochemistry analysis of paraffin-embedded kidney tissues with DAB staining using HXK I Monoclonal Antibody.



Immunofluorescence analysis of NIH/3T3 cells using HXK I Monoclonal Antibody (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.



Flow cytometric analysis of K562 cells using HXK I Monoclonal Antibody (green) and negative control (purple).