



AMPK β 2 Polyclonal Antibody

Catalog No	YP-Ab-14287
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	PRKAB2
Protein Name	5'-AMP-activated protein kinase subunit beta-2
Immunogen	Recombinant Protein of AMPK β 2
Specificity	The antibody detects endogenous AMPK β 2 protein.
Formulation	PBS, pH 7.4, containing 0.5%BSA, 0.02% sodium azide as Preservative and 50% Glycerol.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1:1000-2000
Concentration	1 mg/ml
Purity	$\geq 90\%$
Storage Stability	-20°C/1 year
Synonyms	PRKAB2; 5'-AMP-activated protein kinase subunit beta-2; AMPK subunit beta-2
Observed Band	30kD
Cell Pathway	nucleoplasm,cytosol,nucleotide-activated protein kinase complex,
Tissue Specificity	Liver,Pancreas,
Function	function:AMPK is responsible for the regulation of fatty acid synthesis by phosphorylation of acetyl-CoA carboxylase. Also regulates cholesterol synthesis via phosphorylation and inactivation of hydroxymethylglutaryl-CoA reductase and hormone-sensitive lipase. This is a regulatory subunit, may be a positive regulator of AMPK activity. It may also serve as an adapter molecule for the catalytic alpha-subunit.,PTM:Phosphorylated when associated with the catalytic subunit.,similarity:Belongs to the 5'-AMP-activated protein kinase beta subunit family.,subunit:Heterotrimer of an alpha catalytic subunit, a beta and a gamma non-catalytic regulatory subunits.,
Background	The protein encoded by this gene is a regulatory subunit of the AMP-activated protein kinase (AMPK). AMPK is a heterotrimer consisting of an alpha catalytic subunit, and non-catalytic beta and gamma subunits. AMPK is an important energy-sensing enzyme that monitors cellular energy status. In response to cellular metabolic stresses, AMPK is activated, and thus phosphorylates and



inactivates acetyl-CoA carboxylase (ACC) and beta-hydroxy beta-methylglutaryl-CoA reductase (HMGCR), key enzymes involved in regulating de novo biosynthesis of fatty acid and cholesterol. This subunit may be a positive regulator of AMPK activity. It is highly expressed in skeletal muscle and thus may have tissue-specific roles. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jul 2013],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of 1) 293T, 2) HepG2, 3) Mouse Heart Tissue, 4) Rat Heart Tissue using AMPK β 2 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000