



DGK-α Polyclonal Antibody

Catalog No	YP-Ab-14721
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;IHC;IF;ELISA
Gene Name	DGKA
Protein Name	Diacylglycerol kinase alpha
Immunogen	The antiserum was produced against synthesized peptide derived from human DGKA. AA range:304-353
Specificity	DGK-α Polyclonal Antibody detects endogenous levels of DGK-α protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/20000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	DGKA; DAGK; DAGK1; Diacylglycerol kinase alpha; DAG kinase alpha; 80 kDa diacylglycerol kinase; Diglyceride kinase alpha; DGK-alpha
Observed Band	80kD
Cell Pathway	Cytoplasm, cytosol .
Tissue Specificity	Expressed in lymphocytes.
Function	catalytic activity:ATP + 1,2-diacylglycerol = ADP + 1,2-diacyl-sn-glycerol 3-phosphate.,enzyme regulation:Stimulated by calcium and phosphatidylserine. Phosphorylated by protein kinase C.,function:Upon cell stimulation converts the second messenger diacylglycerol into phosphatidate, initiating the resynthesis of phosphatidylinositols and attenuating protein kinase C activity.,similarity:Belongs to the eukaryotic diacylglycerol kinase family.,similarity:Contains 1 DAGKc domain.,similarity:Contains 2 EF-hand domains.,similarity:Contains 2 phorbol-ester/DAG-type zinc fingers.,subunit:Monomer.,tissue specificity:Lymphocytes and oligodendroglial cells.,
Background	The protein encoded by this gene belongs to the eukaryotic diacylglycerol kinase family. It acts as a modulator that competes with protein kinase C for the second messenger diacylglycerol in intracellular signaling pathways. It also plays an important role in the resynthesis of phosphatidylinositols and phosphorylating diacylglycerol to phosphatidic acid. Alternative splicing occurs at this locus and



four transcript variants encoding the same protein have been identified. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


Products Images




Western Blot analysis of various cells using DGK- α Polyclonal Antibody



Western Blot analysis of Lovo cells using DGK- α Polyclonal Antibody



Western blot analysis of lysates from Jurkat cells, using DGKA Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA, pH9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200(room temperature, 30min).