

(Tel: 400-999-8863 ■ Email:Upingbio.163.com



Jamip2 Polyclonal Antibody

-Ab-14802 man;Rat;Mouse;
man;Rat;Mouse;
R-IHC-IF-FLISA
7,11 IO,11 ,LLIO/1
KMIP2
nus kinase and microtubule-interacting protein 2
e antiserum was produced against synthesized peptide derived from human KMIP2. AA range:761-810
nip2 Polyclonal Antibody detects endogenous levels of Jamip2 protein.
uid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
yclonal, Rabbit,IgG
e antibody was affinity-purified from rabbit antiserum by nity-chromatography using epitope-specific immunogen.
estern Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. munofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other plications.
ng/ml
0%
°C/1 year
KMIP2; JAMIP2; KIAA0555; NECC1; Janus kinase and microtubule-interacting tein 2; CTCL tumor antigen HD-CL-04; Neuroendocrine long coiled-coil protein
KD .
gi apparatus .
hly expressed in brain, moderately expressed in thymus, spleen and lung, and akly expressed in kidney, liver and peripheral blood lymphocytes. Also pressed in adrenal and pituitary glands, as well as testis.
ilarity:Belongs to the JAKMIP family.,tissue specificity:Highly expressed in in, moderately expressed in thymus, spleen and lung, and weakly expressed in ney, liver and peripheral blood lymphocytes. Aslo expressed in adrenal and litary glands, as well as testis.,
ne protein encoded by this gene is reported to be a component of the Golgi trix. It may act as a golgin protein by negatively regulating transit of secretory go and by acting as a structural scaffold of the Golgi. Alternative splicing ults in multiple transcript variants. [provided by RefSeq, Aug 2012],



UpingBio technology Co.,Ltd

C Tel: 400-999-8863 🗷 Email:Upingbio.163.com




matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using JAKMIP2 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from LOVO and A549 cells, using JAKMIP2 Antibody. The lane on the right is blocked with the synthesized peptide.