



PI 3-kinase p85 β Polyclonal Antibody

Catalog No	YP-Ab-14917
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;ELISA
Gene Name	PIK3R2
Protein Name	Phosphatidylinositol 3-kinase regulatory subunit beta
Immunogen	The antiserum was produced against synthesized peptide derived from human PI 3-kinase p85beta. AA range:409-458
Specificity	PI 3-kinase p85 β Polyclonal Antibody detects endogenous levels of PI 3-kinase p85 β protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	$\geq 90\%$
Storage Stability	-20°C/1 year
Synonyms	PIK3R2; Phosphatidylinositol 3-kinase regulatory subunit beta; PI3-kinase regulatory subunit beta; PI3K regulatory subunit beta; PtdIns-3-kinase regulatory subunit beta; Phosphatidylinositol 3-kinase 85 kDa regulatory subunit beta; PI3-kinase
Observed Band	85kD
Cell Pathway	nucleus,cytosol,phosphatidylinositol 3-kinase complex,
Tissue Specificity	Brain,Epithelium,Kidney,Placenta,
Function	function: Binds to activated (phosphorylated) protein-tyrosine kinases, through its SH2 domain, and acts as an adapter, mediating the association of the p110 catalytic unit to the plasma membrane.,similarity: Belongs to the PI3K p85 subunit family.,similarity: Contains 1 Rho-GAP domain.,similarity: Contains 1 SH3 domain.,similarity: Contains 2 SH2 domains.,subunit: Heterodimer of a p110 (catalytic) and a p85 (regulatory) subunits.,
Background	Phosphatidylinositol 3-kinase (PI3K) is a lipid kinase that phosphorylates phosphatidylinositol and similar compounds, creating second messengers important in growth signaling pathways. PI3K functions as a heterodimer of a regulatory and a catalytic subunit. The protein encoded by this gene is a



regulatory component of PI3K. Two transcript variants, one protein coding and the other non-protein coding, have been found for this gene. [provided by RefSeq, Dec 2012],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


Products Images




Western Blot analysis of various cells using PI 3-kinase p85β Polyclonal Antibody diluted at 1:500




Western Blot analysis of 453 cells using PI 3-kinase p85β Polyclonal Antibody diluted at 1:500



Western blot analysis of lysate from Jurkat cells, using PI 3-kinase p85β antibody.



Immunohistochemical analysis of paraffin-embedded human Moderately differentiated hepatocellular carcinoma Antibody was diluted at 1:200(4° overnight).



Immunohistochemical analysis of paraffin-embedded human tonsil Antibody was diluted at 1:200(4° overnight).