



Chromogranin A (ABT037) Mouse mAb

Catalog No	YP-Ab-15090
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	IHC
Gene Name	CHGA
Protein Name	beta Granin;betagranin (N-terminal fragment of chromogranin A);catestatin;CgA;CHG A;Chga;chromofungin;Chromogranin A;Chromogranin A parathyroid secretory protein 1;Chromogranin A precursor;Chromograni
Immunogen	Synthesized peptide derived from human Chromogranin A
Specificity	The antibody can specifically recognize human Chromogranin A protein.
Formulation	PBS, pH7.2, 0.03% Porcolin 300, containing stabilizing protein
Source	Mouse, Monoclonal/IgG1, Kappa
Purification	The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.
Dilution	IHC-p 1:200-400,
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	beta Granin;betagranin (N-terminal fragment of chromogranin A);catestatin;CgA;CHG A;Chga;chromofungin;Chromogranin A;Chromogranin A parathyroid secretory protein 1;Chromogranin A precursor;ChromograninA;CMGA_HUMAN;ER-37;Pancreastatin;Parastatin;Parat hyroid secretory protein 1;Pituitary secretory protein I;Secretory protein I;SP I;SP-I;SP1;SPI;vasostatin 2;Vasostatin;Vasostatin I;Vasostatin II;vasostatin-2
Observed Band	
Cell Pathway	Cytoplasmic
Tissue Specificity	Pancreas/ stomach
Function	function:Pancreastatin strongly inhibits glucose induced insulin release from the pancreas.,miscellaneous:Binds calcium with a low-affinity.,PTM:Sulfated on tyrosine residues and/or contains sulfated glycans.,similarity:Belongs to the chromogranin/secretogranin protein family.,subcellular location:Neuroendocrine and endocrine secretory granules.,
Background	The protein encoded by this gene is a member of the chromogranin/secretogranin family of neuroendocrine secretory proteins. It is



found in secretory vesicles of neurons and endocrine cells. This gene product is a precursor to three biologically active peptides; vasostatin, pancreastatin, and parastatin. These peptides act as autocrine or paracrine negative modulators of the neuroendocrine system. Two other peptides, catestatin and chromofungin, have antimicrobial activity and antifungal activity, respectively. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2014],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


Products Images



Human colon tissue was stained with Anti-Chromogranin A (ABT037) Antibody



Human pancreas tissue was stained with Anti-Chromogranin A (ABT037) Antibody



Human parathyroidoma tissue was stained with Anti-Chromogranin A (ABT037) Antibody