



COX-2 mouse mAb(ABT040)

Catalog No	YP-Ab-15569
Isotype	IgG
Reactivity	Human; Predict react with Mouse, Rat
Applications	IHC;WB;IF
Gene Name	PTGS2 COX2
Protein Name	COX-2
Immunogen	Synthesized peptide derived from human COX-2
Specificity	The antibody can specifically recognize human COX-2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.133% sodium azide.
Source	Mouse, Monoclonal/IgG2b, Kappa
Purification	The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.
Dilution	IHC-p 1:100-500, WB 1:200-1000, IF 1:100-500
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Prostaglandin G/H synthase 2 (EC 1.14.99.1;Cyclooxygenase-2;COX-2;PHS II;Prostaglandin H2 synthase 2;PGH synthase 2;PGHS-2;Prostaglandin-endoperoxide synthase 2)
Observed Band	
Cell Pathway	Microsome membrane ; Peripheral membrane protein. Endoplasmic reticulum membrane ; Peripheral membrane protein. Nucleus inner membrane ; Peripheral membrane protein. Nucleus outer membrane ; Peripheral membrane protein. Detected on the luminal side of the endoplasmic reticulum and nuclear envelope.
Tissue Specificity	Endothelial cell,Epidermal keratinocytes in primary culture,Lung,Pe
Function	catalytic activity:Arachidonate + AH(2) + 2 O(2) = prostaglandin H(2) + A + H(2)O.,cofactor: Binds 1 heme B (iron-protoporphyrin IX) group per subunit.,disease: Likely to play a role in inflammatory diseases such as rheumatoid arthritis.,function: May have a role as a major mediator of inflammation and/or a role for prostanoid signaling in activity-dependent plasticity.,induction: By cytokines and mitogens.,miscellaneous: This enzyme acts both as a dioxygenase and as a peroxidase.,miscellaneous: This enzyme is the target of nonsteroidal anti-inflammatory drugs such as aspirin.,pathway: Lipid metabolism; prostaglandin biosynthesis.,similarity: Belongs to the prostaglandin G/H synthase family.,similarity: Contains 1 EGF-like domain.,subunit: Homodimer.,



Background

Prostaglandin-endoperoxide synthase (PTGS), also known as cyclooxygenase, is the key enzyme in prostaglandin biosynthesis, and acts both as a dioxygenase and as a peroxidase. There are two isozymes of PTGS: a constitutive PTGS1 and an inducible PTGS2, which differ in their regulation of expression and tissue distribution. This gene encodes the inducible isozyme. It is regulated by specific stimulatory events, suggesting that it is responsible for the prostanoid biosynthesis involved in inflammation and mitogenesis. [provided by RefSeq, Feb 2009],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Human colon carcinoma tissue was stained with Anti-COX-2 (ABT040) Antibody



Hela whole cell lysates were separated by 10% SDS-PAGE, and the membrane was blotted with anti-COX-2(ABT040) antibody. The HRP-conjugated Goat anti-Mouse IgG(H + L) antibody was used to detect the antibody. Lane 1: HeLa Predicted band size: 70kDa Observed band size: 70kDa