



GRO α Polyclonal Antibody

Catalog No	YP-Ab-15921
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	IHC;IF;ELISA
Gene Name	CXCL1
Protein Name	Growth-regulated alpha protein
Immunogen	The antiserum was produced against synthesized peptide derived from human GRO α . AA range:39-88
Specificity	GRO α Polyclonal Antibody detects endogenous levels of GRO α protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	IHC: 1/100 - 1/300. ELISA: 1/10000.. IF 1:50-200
Concentration	1 mg/ml
Purity	$\geq 90\%$
Storage Stability	-20°C/1 year
Synonyms	CXCL1; GRO; GRO1; GROA; MGSA; SCYB1; Growth-regulated alpha protein; C-X-C motif chemokine 1; GRO-alpha(1-73); Melanoma growth stimulatory activity; MGSA; Neutrophil-activating protein 3; NAP-3
Observed Band	
Cell Pathway	Secreted.
Tissue Specificity	Blood,Ovary,Peripheral blood monocyte,Placenta,Skin,
Function	function:Has chemotactic activity for neutrophils. May play a role in inflammation and exerts its effects on endothelial cells in an autocrine fashion. In vitro, the processed forms GRO-alpha(4-73), GRO-alpha(5-73) and GRO-alpha(6-73) show a 30-fold higher chemotactic activity.,online information:CXCL1 entry,PTM:N-terminal processed forms GRO-alpha(4-73), GRO-alpha(5-73) and GRO-alpha(6-73) are produced by proteolytic cleavage after secretion from peripheral blood monocytes.,similarity:Belongs to the intercrine alpha (chemokine CxC) family.,
Background	This antimicrobial gene encodes a member of the CXC subfamily of chemokines. The encoded protein is a secreted growth factor that signals through the G-protein coupled receptor, CXC receptor 2. This protein plays a role in inflammation and as a chemoattractant for neutrophils. Aberrant expression of this protein is



associated with the growth and progression of certain tumors. A naturally occurring processed form of this protein has increased chemotactic activity. Alternate splicing results in coding and non-coding variants of this gene. A pseudogene of this gene is found on chromosome 4. [provided by RefSeq, Sep 2014],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemistry analysis of GRO α antibody in paraffin-embedded human brain tissue.