

C Tel: 400-999-8863 🗷 Email:Upingbio.163.com





## TNF-IP 8 Polyclonal Antibody

Catalog No	YP-Ab-15970
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;IHC;IF;ELISA
Gene Name	TNFAIP8
Protein Name	Tumor necrosis factor alpha-induced protein 8
Immunogen	The antiserum was produced against synthesized peptide derived from human TFIP8. AA range:31-80
Specificity	TNF-IP 8 Polyclonal Antibody detects endogenous levels of TNF-IP 8 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms  Observed Band	TNFAIP8; Tumor necrosis factor alpha-induced protein 8; TNF alpha-induced protein 8; Head and neck tumor and metastasis-related protein; MDC-3.13; NF-kappa-B-inducible DED-containing protein; NDED; SCC-S2; TNF-induced protein GG2-1 23kD
Observed Band	
Cell Pathway	Cytoplasm .
Tissue Specificity	Expressed at high levels in the spleen, lymph node, thymus, thyroid, bone marrow and placenta. Expressed at high levels both in various tumor tissues, unstimulated and cytokine-activated cultured cells. Expressed at low levels in the spinal cord, ovary, lung, adrenal glands, heart, brain, testis and skeletal muscle.
Function	developmental stage:Expressed at high levels in the fetal liver, lung and kidney.,function:Acts as a negative mediator of apoptosis and may play a role in tumor progression. Suppresses the TNF-mediated apoptosis by inhibiting caspase-8 activity but not the processing of procaspase-8, subsequently resulting in inhibition of BID cleavage and caspase-3 activation.,induction:By nuclear factor-KB (NF-KB) and TNF. Induction by TNF depends upon activation of NF-KB.,similarity:Belongs to the TNFAIP8 family.,tissue specificity:Expressed at high levels in the spleen, lymph node, thymus, thyroid, bone marrow and




## UpingBio technology Co.,Ltd

📞 Tel: 400-999-8863 🗷 Email:Upingbio.163.com




placenta. Expressed at high levels both in various tumor tissues, unstimulated and cytokine-activated cultured cells. Expressed at low levels in the spinal cord, ovary, lung, adrenal glands, heart, brain, testis and skeletal muscle., Background developmental stage: Expressed at high levels in the fetal liver, lung and kidney.,function:Acts as a negative mediator of apoptosis and may play a role in tumor progression. Suppresses the TNF-mediated apoptosis by inhibiting caspase-8 activity but not the processing of procaspase-8, subsequently resulting in inhibition of BID cleavage and caspase-3 activation.,induction:By nuclear factor-KB (NF-KB) and TNF. Induction by TNF depends upon activation of NF-KB.,similarity:Belongs to the TNFAIP8 family.,tissue specificity:Expressed at high levels both in various tumor tissues, unstimulated and cytokine-activated cultured cells. Expressed at low levels in the spinal cord, overversely. cytokine-activated cultured cells. Expressed at low levels in the spinal cord, ovary, lung, adrenal glands, heart, brain, testis and skeletal muscle., Avoid repeated freezing and thawing! matters needing attention **Usage suggestions** This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


## **Products Images**



Western Blot analysis of various cells using TNF-IP 8 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunofluorescence analysis of A549 cells, using TFIP8 Antibody. The picture on the right is blocked with the synthesized peptide.




Immunohistochemistry analysis of paraffin-embedded human lung carcinoma tissue, using TFIP8 Antibody. The picture on the right is blocked with the synthesized peptide.



## UpingBio technology Co.,Ltd

**(** Tel: 400-999-8863 **(** Emall:Upingbio.163.com





Western blot analysis of lysates from HUVEC cells, using TFIP8 Antibody. The lane on the right is blocked with the synthesized peptide.