







## PHF13 rabbit pAb


Catalog No	YP-Ab-12280
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB;IHC
Gene Name	PHF13
Protein Name	PHF13
Immunogen	Synthesized peptide derived from human PHF13 AA range: 206-256
Specificity	This antibody detects endogenous levels of PHF13 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20°C/1 year
	-20°C/1 year
Synonyms	-20°C/1 year  Nucleus . Nucleus, nucleoplasm . Predominantly bound to chromatin, but a minor proportion is also detected in the nucleoplasm.
Synonyms Observed Band	Nucleus . Nucleus, nucleoplasm . Predominantly bound to chromatin, but a minor
Synonyms Observed Band Cell Pathway	Nucleus . Nucleus, nucleoplasm . Predominantly bound to chromatin, but a minor
Synonyms Observed Band Cell Pathway Tissue Specificity	Nucleus . Nucleus, nucleoplasm . Predominantly bound to chromatin, but a minor proportion is also detected in the nucleoplasm.
Synonyms  Observed Band  Cell Pathway  Tissue Specificity  Function	Nucleus . Nucleus, nucleoplasm . Predominantly bound to chromatin, but a minor proportion is also detected in the nucleoplasm.
Synonyms  Observed Band  Cell Pathway  Tissue Specificity  Function  Background  matters needing	Nucleus . Nucleus, nucleoplasm . Predominantly bound to chromatin, but a minor proportion is also detected in the nucleoplasm.  similarity:Contains 1 PHD-type zinc finger.,








## **Products Images**



Western blot analysis of lysates from U2OS cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).