







PGBD4 rabbit pAb

Trichonlusia ni. This family also includes genes in several genomes, including		
Reactivity Human;Rat;Mouse; Applications WB Gene Name PGBD4 Protein Name PGBD4 Immunogen Synthesized peptide derived from human PGBD4 AA range: 23-73 Specificity This antibody detects endogenous levels of PGBD4 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Function Background The piggyBac family of proteins, found in diverse animals, are transposases related to the transposase of the canonical piggys act ransposon from the month.	Catalog No	YP-Ab-12304
Applications Gene Name PGBD4 Protein Name PGBD4 Immunogen Synthesized peptide derived from human PGBD4 AA range: 23-73 Specificity This antibody detects endogenous levels of PGBD4 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Function Background The piggyBac family of proteins, found in diverse animals, are transposases related to the transposase of the canonical piggyBac transposan from the manual production.	Isotype	IgG
Gene Name PGBD4 Protein Name PGBD4 Immunogen Synthesized peptide derived from human PGBD4 AA range: 23-73 Specificity This antibody detects endogenous levels of PGBD4 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Function Background The piggyBac family of proteins, found in diverse animals, are transposases related to the transposase of the canonical piggyBac transposan from the Drichopusia in This family as includes genes in several genomes including	Reactivity	Human;Rat;Mouse;
Protein Name PGBD4 Immunogen Synthesized peptide derived from human PGBD4 AA range: 23-73 Specificity This antibody detects endogenous levels of PGBD4 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Function Background The piggyBac family of proteins, found in diverse animals, are transposases related to the transposase of the canonical piggyBac transposon from the moth Trichoplusia in Tis family also includes genes in several penomes including.	Applications	WB
Immunogen Synthesized peptide derived from human PGBD4 AA range: 23-73 Specificity This antibody detects endogenous levels of PGBD4 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Function Background The piggyBac family of proteins, found in diverse animals, are transposases related to the transposase of the canonical piggyBac transposon from the moth Trichoplusia in 1 This family also includes genes in several grappes in several grappes in several grappes.	Gene Name	PGBD4
Specificity This antibody detects endogenous levels of PGBD4 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Function Background The piggyBac family of proteins, found in diverse animals, are transposases related to the transposase of the canonical piggyBac transposon from the month of transposase of the canonical piggyBac transposon from the printy and the proteins are transposases including.	Protein Name	PGBD4
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Function Background The piggyBac family of proteins, found in diverse animals, are transposases related to the transposase of the canonical piggyBac transposant from the first family also includes genes in several genomes, including includes genes in several genomes including	Immunogen	Synthesized peptide derived from human PGBD4 AA range: 23-73
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Function Background The piggyBac family of proteins, found in diverse animals, are transposases related to the transposase of the canonical piggyBac transposon from the moth Trichon lusia ni This family also includes genes in several genomes including	Specificity	This antibody detects endogenous levels of PGBD4 at Human
Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Function Background The piggyBac family of proteins, found in diverse animals, are transposases related to the transposase of the canonical piggyBac transposon from the more transposase of the canonical piggyBac transposon from the more transposase of the canonical piggyBac transposon from the more transposase of the canonical piggyBac transposon from the more transposase of the canonical piggyBac transposon from the more transposase of the canonical piggyBac transposon from the more transposase of the canonical piggyBac transposon from the more transposase of the canonical piggyBac transposon from the more transposase of the canonical piggyBac transposon from the more transposase of the canonical piggyBac transposon from the more transposase of the canonical piggyBac transposon from the more transposase of the canonical piggyBac transposon from the more transposase of the canonical piggyBac transposon from the more transposase of the canonical piggyBac transposon from the more transposase of the canonical piggyBac transposon from the more transposase of the canonical piggyBac transposon from the more transposase of the canonical piggyBac transposon from the more transposase of the canonical piggyBac transposon from the more transposase of the canonical piggyBac transposon from the more transposase of the piggyBac transp	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Function Background The piggyBac family of proteins, found in diverse animals, are transposases related to the transposase of the canonical piggyBac transposanes in several genome the moth Trichonylusia in This family also indudes genes in several genomes including	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Function Background The piggyBac family of proteins, found in diverse animals, are transposases related to the transposase of the canonical piggyBac transposon from the moth Trichpollusia ni This family also includes genes in several genomes including	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Function Background The piggyBac family of proteins, found in diverse animals, are transposases related to the transposase of the canonical piggyBac transposon from the moth Trichonlusia ni. This family also includes genes in several genomes, including	Dilution	WB 1: 500-2000
Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Function Background The piggyBac family of proteins, found in diverse animals, are transposases related to the transposase of the canonical piggyBac transposon from the moth Trichoplusia ni. This family also includes genes in several genomes, including	Concentration	1 mg/ml
Synonyms Observed Band Cell Pathway Tissue Specificity Function Background The piggyBac family of proteins, found in diverse animals, are transposases related to the transposase of the canonical piggyBac transposon from the moth Trichoplusia ni. This family also includes genes in several genomes, including	Purity	≥90%
Observed Band Cell Pathway Tissue Specificity Function Background The piggyBac family of proteins, found in diverse animals, are transposases related to the transposase of the canonical piggyBac transposon from the moth Trichonlusia ni. This family also includes genes in several genomes, including	Storage Stability	-20°C/1 year
Cell Pathway Tissue Specificity Function Background The piggyBac family of proteins, found in diverse animals, are transposases related to the transposase of the canonical piggyBac transposon from the moth Trichoplusia ni. This family also includes genes in several genomes, including	Synonyms	
Tissue Specificity Function Background The piggyBac family of proteins, found in diverse animals, are transposases related to the transposase of the canonical piggyBac transposon from the moth Trichoplusia ni. This family also includes genes in several genomes, including	Observed Band	
Function Background The piggyBac family of proteins, found in diverse animals, are transposases related to the transposase of the canonical piggyBac transposon from the moth Trichoplusia ni. This family also includes genes in several genomes, including	Cell Pathway	
Background The piggyBac family of proteins, found in diverse animals, are transposases related to the transposase of the canonical piggyBac transposon from the moth Trichoplusia ni. This family also includes genes in several genomes, including	Tissue Specificity	
related to the transposase of the canonical piggyBac transposon from the moth	Function	
genes. The PGBD proteins appear to be novel, with no obvious relationship to other transposases, or other known protein families. [provided by RefSeq, Jul 2008],	Background	related to the transposase of the canonical piggyBac transposon from the moth, Trichoplusia ni. This family also includes genes in several genomes, including human, that appear to have been derived from the piggyBac transposons. This gene belongs to the subfamily of piggyBac transposable element derived (PGBD genes. The PGBD proteins appear to be novel, with no obvious relationship to other transposases, or other known protein families. [provided by RefSeq, Jul
matters needing Avoid repeated freezing and thawing! attention		Avoid repeated freezing and thawing!



UpingBio technology Co.,Ltd


C Tel: 400-999-8863 ■ Email:UpingBio@163.com



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from MCF-7 cells, primary antibody was diluted at 1:1000, 4° over night