







LARP4 rabbit pAb

| Catalog No | YP-Ab-12309 |
|---|---|
| Isotype | IgG |
| Reactivity | Human; Mouse |
| Applications | WB;ELISA;IHC |
| Gene Name | LARP4 PP13296 |
| Protein Name | LARP4 |
| Immunogen | Synthesized peptide derived from human LARP4 AA range: 209-259 |
| Specificity | This antibody detects endogenous levels of LARP4 at Human/Mouse |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. |
| Dilution | WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000 |
| | |
| Concentration | 1 mg/ml |
| Concentration Purity | 1 mg/ml ≥90% |
| | • |
| Purity | ≥90% |
| Purity Storage Stability | ≥90% |
| Purity Storage Stability Synonyms | ≥90% |
| Purity Storage Stability Synonyms Observed Band | ≥90% -20°C/1 year Cytoplasm, Stress granule . Cytoplasm, cytosol . Localized throughout the |
| Purity Storage Stability Synonyms Observed Band Cell Pathway | ≥90% -20°C/1 year Cytoplasm, Stress granule . Cytoplasm, cytosol . Localized throughout the |
| Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity | ≥90% -20°C/1 year Cytoplasm, Stress granule . Cytoplasm, cytosol . Localized throughout the cytosol. Partially localized in stress granules in response to arsenite treatment sequence caution:Translation N-terminally extended.,similarity:Contains 1 HTH La-type RNA-binding domain.,similarity:Contains 1 RRM (RNA recognition motif) |
| Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function | ≥90% -20°C/1 year Cytoplasm, Stress granule . Cytoplasm, cytosol . Localized throughout the cytosol. Partially localized in stress granules in response to arsenite treatment sequence caution:Translation N-terminally extended.,similarity:Contains 1 HTH La-type RNA-binding domain.,similarity:Contains 1 RRM (RNA recognition motif) |
| Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing | ≥90% -20°C/1 year Cytoplasm, Stress granule . Cytoplasm, cytosol . Localized throughout the cytosol. Partially localized in stress granules in response to arsenite treatment sequence caution:Translation N-terminally extended.,similarity:Contains 1 HTH La-type RNA-binding domain.,similarity:Contains 1 RRM (RNA recognition motif) domain., |

Products Images



Western blot analysis of lysates from K562 cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).