





AGAP9 rabbit pAb

Isotype IgG Reactivity Human;Rat;Mouse; Applications WB Gene Name AGAP9 CTGLF6 Protein Name AGAP9 Immunogen Synthesized peptide derived from human AGAP9 AA range: 4-54 Specificity This antibody detects endogenous levels of AGAP9 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Function caution: The sequence shown here is derived from an Ensembliautomatic analysis		
Reactivity Human;Rat;Mouse; Applications WB Gene Name AGAP9 CTGLF6 Protein Name AGAP9 Immunogen Synthesized peptide derived from human AGAP9 AA range: 4-54 Specificity This antibody detects endogenous levels of AGAP9 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity	Catalog No	YP-Ab-12342
Applications WB Gene Name AGAP9 CTGLF6 Protein Name AGAP9 Immunogen Synthesized peptide derived from human AGAP9 AA range: 4-54 Specificity This antibody detects endogenous levels of AGAP9 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity	Isotype	IgG
Gene Name AGAP9 CTGLF6 Protein Name AGAP9 Immunogen Synthesized peptide derived from human AGAP9 AA range: 4-54 Specificity This antibody detects endogenous levels of AGAP9 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity	Reactivity	Human;Rat;Mouse;
Protein Name AGAP9 Immunogen Synthesized peptide derived from human AGAP9 AA range: 4-54 Specificity This antibody detects endogenous levels of AGAP9 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, lgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity	Applications	WB
Immunogen Synthesized peptide derived from human AGAP9 AA range: 4-54 Specificity This antibody detects endogenous levels of AGAP9 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity	Gene Name	AGAP9 CTGLF6
Specificity This antibody detects endogenous levels of AGAP9 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity	Protein Name	AGAP9
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity	Immunogen	Synthesized peptide derived from human AGAP9 AA range: 4-54
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity	Specificity	This antibody detects endogenous levels of AGAP9 at Human
Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity	Dilution	WB 1: 500-2000
Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity	Concentration	1 mg/ml
Synonyms Observed Band Cell Pathway Tissue Specificity	Purity	≥90%
Observed Band Cell Pathway Tissue Specificity	Storage Stability	-20°C/1 year
Cell Pathway Tissue Specificity	Synonyms	
Tissue Specificity	Observed Band	
	Cell Pathway	
Function caution: The sequence shown here is derived from an Ensembl automatic analysis	Tissue Specificity	
pipeline and should be considered as preliminary data.,function:Putative GTPase-activating protein .,miscellaneous:Encoded by one of the numerous	Function	GTPase-activating protein .,miscellaneous:Encoded by one of the numerous copies of centaurin gamma-like genes clustered in the q11 region of chromosome 10.,similarity:Belongs to the centaurin gamma-like family.,similarity:Contains 1 ANK repeat.,similarity:Contains 1 Arf-GAP domain.,similarity:Contains 1 PH
Background	Background	
matters needing Avoid repeated freezing and thawing! attention	matters needing attention	Avoid repeated freezing and thawing!



UpingBio technology Co.,Ltd


C Tel: 400-999-8863 ■ Email:UpingBio@163.com



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from MCF-7 cells, primary antibody was diluted at 1:1000, 4° over night