






## UNKL rabbit pAb

| Catalog No                | YP-Ab-12346   |
|---------------------------|---|
| Isotype                   | IgG   |
| Reactivity                | Human; Mouse  |
| Applications              | WB  |
| Gene Name                 | UNKL C16orf28 ZC3H5L ZC3HDC5L   |
| Protein Name              | UNKL  |
| Immunogen                 | Synthesized peptide derived from human UNKL AA range: 296-346   |
| Specificity               | This antibody detects endogenous levels of UNKL at Human/Mouse  |
| Formulation               | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| Source                    | Polyclonal, Rabbit,IgG  |
| Purification              | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.   |
| Dilution                  | WB 1: 500-2000  |
| Concentration             | 1 mg/ml   |
| Purity                    | ≥90%  |
| Storage Stability         | -20°C/1 year  |
| Synonyms                  |   |
| Observed Band             |   |
| Cell Pathway              | [Isoform 4]: Cytoplasm. Nucleus. Isoform 4 is primarily localized in the cytoplasm but has the ability to shuttle between the nucleus and the cytoplasm.  |
| Tissue Specificity        |   |
| Function                  | similarity:Belongs to the unkempt family.,similarity:Contains 1 RING-type zinc finger.,   |
| Background                | This gene encodes a RING finger protein that may function in Rac signaling. It can bind to Brg/Brm-associated factor 60b and can promote its ubiquitination. Alternative splicing of this gene results in multiple transcript variants. [provided by RefSeq, Jun 2013], |
| matters needing attention | Avoid repeated freezing and thawing!  |
| Usage suggestions         | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.   |
|                           |   |



## **Products Images**



Western blot analysis of lysates from KB cells, primary antibody was diluted at 1:1000, 4°over night