

SNAT rabbit pAb

Catalog No	YP-Ab-12511
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;IHC
Gene Name	AANAT SNAT
Protein Name	SNAT
Immunogen	Synthesized peptide derived from human SNAT
Specificity	This antibody detects endogenous levels of Human, Mouse, Rat SNAT
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
	-20 O/T year
Synonyms	Serotonin N-acetyltransferase (Serotonin acetylase;EC 2.3.1.87;Aralkylamine N-acetyltransferase;AA-NAT)
Synonyms Observed Band	Serotonin N-acetyltransferase (Serotonin acetylase;EC 2.3.1.87;Aralkylamine
-	Serotonin N-acetyltransferase (Serotonin acetylase;EC 2.3.1.87;Aralkylamine
Observed Band	Serotonin N-acetyltransferase (Serotonin acetylase;EC 2.3.1.87;Aralkylamine N-acetyltransferase;AA-NAT)
Observed Band Cell Pathway	Serotonin N-acetyltransferase (Serotonin acetylase;EC 2.3.1.87;Aralkylamine N-acetyltransferase;AA-NAT) Cytoplasm . Highly expressed in pineal gland and at lower levels in the retina. Weak
Observed Band Cell Pathway Tissue Specificity	Serotonin N-acetyltransferase (Serotonin acetylase;EC 2.3.1.87;Aralkylamine N-acetyltransferase;AA-NAT) Cytoplasm . Highly expressed in pineal gland and at lower levels in the retina. Weak
Observed Band Cell Pathway Tissue Specificity Function	Serotonin N-acetyltransferase (Serotonin acetylase;EC 2.3.1.87;Aralkylamine N-acetyltransferase;AA-NAT) Cytoplasm . Highly expressed in pineal gland and at lower levels in the retina. Weak
Observed Band Cell Pathway Tissue Specificity Function Background matters needing	Serotonin N-acetyltransferase (Serotonin acetylase;EC 2.3.1.87;Aralkylamine N-acetyltransferase;AA-NAT) Cytoplasm . Highly expressed in pineal gland and at lower levels in the retina. Weak expression in several brain regions and in the pituitary gland.




UpingBio technology Co.,Ltd







Products Images



Immunohistochemical analysis of paraffin-embedded human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200(room temperature, 45min).