





GAP-43 (phospho Ser41) Polyclonal Antibody

sidé . Cell junction, synapse . Cell projection, filopodium membrane ; Peripheral		
Reactivity Human;Mouse;Rat Applications IF;ELISA Gene Name GAP43 Protein Name Neuromodulin Immunogen The antiserum was produced against synthesized peptide derived from human GAP43 around the phosphorylation site of Ser41. AA range;8-57 Specificity Phospho-GAP-43 (S41) Polyclonal Antibody detects endogenous levels of GAP-43 protein only when phosphorylated at S41. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms GAP43; Neuromodulin; Axonal membrane protein GAP-43; Growth-associated protein 43; Neural phosphoprotein B-50; pp46 Observed Band Cell Pathway Cell membrane; Peripheral membrane protein; Cytoplasmic side . Cell projection, growth cone membrane; Peripheral membrane protein; Cytoplasmic side . Cell projection, dendrite. Cell projection, axon Cytoplasm cytoplasmic surface of growth cone and synaptic plasma membranes. Tissue Specificity Alzheimer cortex, Brain, Subthalamic nucleus, function: This protein is associated with nerve growth. It is a major component of the motile "growth cones" that form the tips of elongating axons, online information: Gap-43 entry, PTM:-Phosphorylation of this protein by a protein kinas C is specifically correlated with certain forms of synaptic plasmin membrane is protein in protein plasma membrane of the motile "growth cones" that form the tips of elongating axons, online information: Gap-43 entry, PTM:-Phosphorylation of this protein by a protein kinas C is specifically correlated with certain forms of synaptic plasma membranes.	Catalog No	YP-Ab-12634
Applications IF,ELISA Gene Name GAP43 Protein Name Neuromodulin Immunogen The antiserum was produced against synthesized peptide derived from human GAP43 around the phosphorylation site of Ser41. AA range:8-57 Specificity Phospho-GAP-43 (S41) Polyclonal Antibody detects endogenous levels of GAP-43 protein only when phosphorylated at S41. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms GAP43; Neuromodulin; Axonal membrane protein GAP-43; Growth-associated protein 43; Neural phosphoprotein B-50; pp46 Observed Band Cell Pathway Cell projection, growth cone membrane; Peripheral membrane protein; Cytoplasmic side. Cell junction, synapse. Cell projection, filopodium membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection, endrite. Cell projection, axon Cytoplasmic surface of growth cone and synaptic plasma membranes. Tissue Specificity Alzheimer cortex, Brain, Subthalamic nucleus, function: This protein is associated with nerve growth. It is a major component of the motile "growth cones" that form the tips of elongating axons, online information: Gap-43 estiny. PTM: Phosphorylation of this protein by a protein kinas C is specifically correlated with certain forms of synaptic plasma membrane membrane membrane membranes usubunite. Binds calmodulin family, similarity: Contains 1 IQ domain, subcellular location: Cytoplasmic surface of growth cone and synaptic plasma membranes amembranes assubunite. Binds calmodulin with a greater affinity in the	Isotype	IgG
Gene Name GAP43	Reactivity	Human;Mouse;Rat
Protein Name Neuromodulin	Applications	IF;ELISA
Immunogen The antiserum was produced against synthesized peptide derived from human GAP43 around the phosphorylation site of Ser41. AA range:8-57 Specificity Phospho-GAP-43 (S41) Polyclonal Antibody detects endogenous levels of GAP-43 protein only when phosphorylated at S41. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms GAP43; Neuromodulin; Axonal membrane protein GAP-43; Growth-associated protein 43; Neural phosphoprotein B-50; pp46 Observed Band Cell Pathway Cell projection, growth cone membrane: Peripheral membrane protein: Cytoplasmiside. Cell projection, synapse. Cell projection, fliopodium membrane: Peripheral membrane protein: Perikanyon. Cell projection, dendrite. Cell projection, axon Cytoplasm: Cytoplasmic surface of growth cone and synaptic plasma membranes. Tissue Specificity Alzheimer cortex,Brain,Subthalamic nucleus, Function function:This protein is associated with nerve growth. It is a major component of the motile "growth cones" t	Gene Name	GAP43
GAP43 around the phosphorylation site of Ser41. AA range:8-57 Specificity Phospho-GAP-43 (S41) Polyclonal Antibody detects endogenous levels of GAP-43 protein only when phosphorylated at S41. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms GAP43; Neuromodulin; Axonal membrane protein GAP-43; Growth-associated protein 43; Neural phosphoprotein B-50; pp46 Observed Band Cell Pathway Cell membrane ; Peripheral membrane protein ; Cytoplasmic side. Cell projection, growth cone membrane ; Peripheral membrane protein ; Cytoplasmic side. Cell projection, growth cone membrane ; Peripheral membrane protein ; Cytoplasmic side. Cell projection, dendrite. Cell projection, axon Cytoplasm. Cytoplasmic surface of growth cone and synaptic plasma membranes. Tissue Specificity Alzheimer cortex, Brain, Subthalamic nucleus, Function function: This protein is associated with nerve growth. It is a major component of the motile "growth cones" that form the tips of elongating axons., online information: Gap-43 entry, PTM:Phosphorylation of this protein by a protein kinas C is specifically correlated with certain forms of synaptic plasmia membranes, subunit:Binds calmodulin with a greater affinity in the	Protein Name	Neuromodulin
GAP-43 protein only when phosphorylated at S41. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms GAP43; Neuromodulin; Axonal membrane protein GAP-43; Growth-associated protein 43; Neural phosphoprotein B-50; pp46 Observed Band Cell Pathway Cell membrane ; Peripheral membrane protein ; Cytoplasmic side . Cell projection, growth cone membrane ; Peripheral membrane protein ; Cytoplasmic side . Cell projection, growth cone membrane of growth cone and synaptic plasma membranes. Tissue Specificity Alzheimer cortex, Brain, Subthalamic nucleus, Function function: This protein is associated with nerve growth. It is a major component of the motile "growth cones" that form the tips of elongating axons. online information: Gap-43 entry, PTM:Phosphorylation of this protein by a protein kinas C is specifically correlated with certain forms of synaptic plasticity, similarity:Belongs to the neuromodulin family, similarity:Contains 1 IQ domain, subcellular location:Cytoplasmic surface of growth cone are affinity in the	Immunogen	
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms GAP43; Neuromodulin; Axonal membrane protein GAP-43; Growth-associated protein 43; Neural phosphoprotein B-50; pp46 Observed Band Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection, growth cone membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection, synapse. Cell projection, filopodium membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection, dendrite. Cell projection, axon Cytoplasmic Surface of growth cone and synaptic plasma membranes. Tissue Specificity Alzheimer cortex,Brain,Subthalamic nucleus, Function function:This protein is associated with nerve growth. It is a major component of the motile "growth cones" that form the tips of elongating axons.,online information:Gap-43 entry,PTM:Phosphorylation of this protein by a protein kinas C is specifically correlated with certain forms of synaptic plasticity, similarity:Belongs to the neuromodulin family, similarity:Contains 1 IQ domain.,subcellular location:Cytoplasmic surface of growth cone and synaptic plasma membranes.,subunit:Binds calmodulin with a greater affinity in the	Specificity	Phospho-GAP-43 (S41) Polyclonal Antibody detects endogenous levels of GAP-43 protein only when phosphorylated at S41.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms GAP43; Neuromodulin; Axonal membrane protein GAP-43; Growth-associated protein 43; Neural phosphoprotein B-50; pp46 Observed Band Cell Pathway Cell membrane ; Peripheral membrane protein ; Cytoplasmic side . Cell projection, growth cone membrane ; Peripheral membrane protein ; Cytoplasmic side . Cell junction, synapse. Cell projection, filopodium membrane : Peripheral membrane protein ; Cytoplasmic side . Cell projection, axon Cytoplasm . Cytoplasmic surface of growth cone and synaptic plasma membranes. Cytoplasmic surface of growth cone and synaptic plasma function: This protein is associated with nerve growth. It is a major component of the motile "growth cones" that form the tips of elongating axons.,online information: Gap-43 entry. PTM:Phosphorylation of this protein by a protein kinas C is specifically correlated with certain forms synaptic plasticity, similarity:Belongs to the neuromodulin family, similarity:Contains 1 IQ domain., subcellular location:Cytoplasmic surface of growth cone and synaptic plasma membranes, subunit:Binds calmodulin with a greater affinity in the	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms GAP43; Neuromodulin; Axonal membrane protein GAP-43; Growth-associated protein 43; Neural phosphoprotein B-50; pp46 Observed Band Cell Pathway Cell membrane ; Peripheral membrane protein ; Cytoplasmic side . Cell projection, growth cone membrane ; Peripheral membrane protein ; Cytoplasmic side . Cell junction, synapse . Cell projection, flopodium membrane ; Peripheral membrane protein . Cell projection, axon Cytoplasm . Cytoplasmic surface of growth cone and synaptic plasma membranes Tissue Specificity Alzheimer cortex,Brain,Subthalamic nucleus, Function function:This protein is associated with nerve growth. It is a major component of the motile "growth cones" that form the tips of elongating axons.,online information:Gap-43 entry, PTM:Phosphorylation of this protein by a protein kinas C is specifically correlated with certain forms of synaptic plasticity.,similarity:Belongs to the neuromodulin family,.similarity:Contains 1 IQ domain,.subcellular location:Cytoplasmic surface of growth cone and synaptic plasma membranes,.subunit:Binds calmodulin with a greater affinity in the	Source	Polyclonal, Rabbit,IgG
applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms GAP43; Neuromodulin; Axonal membrane protein GAP-43; Growth-associated protein 43; Neural phosphoprotein B-50; pp46 Observed Band Cell Pathway Cell membrane ; Peripheral membrane protein ; Cytoplasmic side . Cell projection, growth cone membrane ; Peripheral membrane protein ; Cytoplasmic side . Cell junction, synapse . Cell projection, filopodium membrane ; Peripheral membrane protein . Perikaryon . Cell projection, dendrite . Cell projection, axon Cytoplasm . Cytoplasmic surface of growth cone and synaptic plasma membranes. Tissue Specificity Alzheimer cortex,Brain,Subthalamic nucleus, Function function:This protein is associated with nerve growth. It is a major component of the motile "growth cones" that form the tips of elongating axons.,online information:Gap-43 entry,PTM:Phosphorylation of this protein by a protein kinas C is specifically correlated with certain forms of synaptic plasticity, similarity:Belongs to the neuromodulin family, similarity:Contains 1 IQ domain., subcellular location:Cytoplasmic surface of growth cone and synaptic plasma membranes., subunit:Binds calmodulin with a greater affinity in the	Purification	, , , , , , , , , , , , , , , , , , , ,
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms GAP43; Neuromodulin; Axonal membrane protein GAP-43; Growth-associated protein 43; Neural phosphoprotein B-50; pp46 Observed Band Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection, growth cone membrane; Peripheral membrane protein; Cytoplasmic side. Cell junction, synapse. Cell projection, filopodium membrane; Peripheral membrane; Peripheral membrane; Peripheral membrane; Cell projection, axon Cytoplasm. Cytoplasmic surface of growth cone and synaptic plasma membranes. Tissue Specificity Alzheimer cortex,Brain,Subthalamic nucleus, Function function:This protein is associated with nerve growth. It is a major component of the motile "growth cones" that form the tips of elongating axons, online information:Gap-43 entry,PTM:Phosphorylation of this protein by a protein kinas C is specifically correlated with certain forms of synaptic plasticity, similarity:Belongs to the neuromodulin family, similarity:Contains 1 IQ domain., subcellular location:Cytoplasmic surface of growth cone and synaptic plasma membranes, subunit:Binds calmodulin with a greater affinity in the	Dilution	
Synonyms GAP43; Neuromodulin; Axonal membrane protein GAP-43; Growth-associated protein 43; Neural phosphoprotein B-50; pp46 Observed Band Cell Pathway Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection, growth cone membrane; Peripheral membrane protein; Cytoplasmic side. Cell junction, synapse. Cell projection, filopodium membrane; Peripheral membrane protein. Peripheral membrane protein. Peripheral membrane protein. Synapse. Cell projection, dendrite. Cell projection, axon Cytoplasm. Cytoplasmic surface of growth cone and synaptic plasma membranes Tissue Specificity Alzheimer cortex, Brain, Subthalamic nucleus, function: This protein is associated with nerve growth. It is a major component of the motile "growth cones" that form the tips of elongating axons., online information: Gap-43 entry, PTM: Phosphorylation of this protein by a protein kinase C is specifically correlated with certain forms of synaptic plasticity, similarity: Belongs to the neuromodulin family., similarity: Contains 1 IQ domain., subcellular location: Cytoplasmic surface of growth cone and synaptic plasma membranes., subunit: Binds calmodulin with a greater affinity in the	Concentration	• •
Synonyms GAP43; Neuromodulin; Axonal membrane protein GAP-43; Growth-associated protein 43; Neural phosphoprotein B-50; pp46 Observed Band Cell Pathway Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection, growth cone membrane; Peripheral membrane protein; Cytoplasmic side. Cell junction, synapse. Cell projection, filopodium membrane; Peripheral membrane protein. Perikaryon. Cell projection, dendrite. Cell projection, axon Cytoplasm. Cytoplasmic surface of growth cone and synaptic plasma membranes. Tissue Specificity Alzheimer cortex,Brain,Subthalamic nucleus, function:This protein is associated with nerve growth. It is a major component of the motile "growth cones" that form the tips of elongating axons.,online information:Gap-43 entry,PTM:Phosphorylation of this protein by a protein kinas C is specifically correlated with certain forms of synaptic plasticity,,similarity:Belongs to the neuromodulin family,.similarity:Contains 1 IQ domain.,subcellular location:Cytoplasmic surface of growth cone and synaptic plasma membranes.,subunit:Binds calmodulin with a greater affinity in the	Purity	≥90%
Observed Band Cell Pathway Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection, growth cone membrane; Peripheral membrane protein; Cytoplasmic side. Cell junction, synapse. Cell projection, filopodium membrane; Peripheral membrane protein. Perikaryon. Cell projection, dendrite. Cell projection, axon Cytoplasm. Cytoplasmic surface of growth cone and synaptic plasma membranes. Tissue Specificity Alzheimer cortex,Brain,Subthalamic nucleus, Function function:This protein is associated with nerve growth. It is a major component of the motile "growth cones" that form the tips of elongating axons.,online information:Gap-43 entry,PTM:Phosphorylation of this protein by a protein kinas C is specifically correlated with certain forms of synaptic plasticity.,similarity:Belongs to the neuromodulin family.,similarity:Contains 1 IQ domain.,subcellular location:Cytoplasmic surface of growth cone and synaptic plasma membranes.,subunit:Binds calmodulin with a greater affinity in the	Storage Stability	-20°C/1 year
Cell Pathway Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection, growth cone membrane; Peripheral membrane protein; Cytoplasmic side. Cell junction, synapse. Cell projection, filopodium membrane; Peripheral membrane protein. Perikaryon. Cell projection, dendrite. Cell projection, axon Cytoplasm. Cytoplasmic surface of growth cone and synaptic plasma membranes. Tissue Specificity Alzheimer cortex,Brain,Subthalamic nucleus, function:This protein is associated with nerve growth. It is a major component of the motile "growth cones" that form the tips of elongating axons.,online information:Gap-43 entry,PTM:Phosphorylation of this protein by a protein kinast C is specifically correlated with certain forms of synaptic plasticity.,similarity:Belongs to the neuromodulin family.,similarity:Contains 1 IQ domain.,subcellular location:Cytoplasmic surface of growth cone and synaptic plasma membranes.,subunit:Binds calmodulin with a greater affinity in the	Synonyms	•
projection, growth cone membrane; Peripheral membrane protein; Cytoplasmic side. Cell junction, synapse. Cell projection, filopodium membrane; Peripheral membrane protein. Perikaryon. Cell projection, dendrite. Cell projection, axon Cytoplasm. Cytoplasmic surface of growth cone and synaptic plasma membranes Tissue Specificity Alzheimer cortex,Brain,Subthalamic nucleus, function:This protein is associated with nerve growth. It is a major component of the motile "growth cones" that form the tips of elongating axons.,online information:Gap-43 entry,PTM:Phosphorylation of this protein by a protein kinas C is specifically correlated with certain forms of synaptic plasticity.,similarity:Belongs to the neuromodulin family.,similarity:Contains 1 IQ domain.,subcellular location:Cytoplasmic surface of growth cone and synaptic plasma membranes.,subunit:Binds calmodulin with a greater affinity in the	Observed Band	
function: This protein is associated with nerve growth. It is a major component of the motile "growth cones" that form the tips of elongating axons.,online information: Gap-43 entry, PTM: Phosphorylation of this protein by a protein kinas C is specifically correlated with certain forms of synaptic plasticity., similarity: Belongs to the neuromodulin family., similarity: Contains 1 IQ domain., subcellular location: Cytoplasmic surface of growth cone and synaptic plasma membranes., subunit: Binds calmodulin with a greater affinity in the	Cell Pathway	projection, growth cone membrane; Peripheral membrane protein; Cytoplasmic side. Cell junction, synapse. Cell projection, filopodium membrane; Peripheral membrane protein. Perikaryon. Cell projection, dendrite. Cell projection, axon. Cytoplasm. Cytoplasmic surface of growth cone and synaptic plasma
the motile "growth cones" that form the tips of elongating axons.,online information:Gap-43 entry,PTM:Phosphorylation of this protein by a protein kinas C is specifically correlated with certain forms of synaptic plasticity.,similarity:Belongs to the neuromodulin family.,similarity:Contains 1 IQ domain.,subcellular location:Cytoplasmic surface of growth cone and synaptic plasma membranes.,subunit:Binds calmodulin with a greater affinity in the	Tissue Specificity	Alzheimer cortex,Brain,Subthalamic nucleus,
` '		function. This waste is in a consisted with warms was the latin a waste account of



UpingBio technology Co.,Ltd

📞 Tel: 400-999-8863 🛎 Email:UpingBio@163.com



Background

The protein encoded by this gene has been termed a 'growth' or 'plasticity' protein because it is expressed at high levels in neuronal growth cones during development and axonal regeneration. This protein is considered a crucial component of an effective regenerative response in the nervous system. Alternatively spliced transcript variants encoding distinct isoforms have been found for this gene. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using GAP43 (Phospho-Ser41) Antibody





Immunofluorescence analysis of MCF-7 cells, using GAP43 (Phospho-Ser41) Antibody. The picture on the right is blocked with the phospho peptide.