



EphA5 Monoclonal Antibody

Catalog No	YP-Ab-12909
Isotype	IgG
Reactivity	Human
Applications	WB;ELISA
Gene Name	EPHA5
Protein Name	Ephrin type-A receptor 5
Immunogen	Purified recombinant fragment of EphA5 (aa620-774) expressed in E. Coli.
Specificity	EphA5 Monoclonal Antibody detects endogenous levels of EphA5 protein.
Formulation	Ascitic fluid containing 0.03% sodium azide, 0.5% BSA, 50% glycerol.
Source	Monoclonal, Mouse
Purification	Affinity purification
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	EPHA5; BSK; EHK1; HEK7; TYRO4; Ephrin type-A receptor 5; Brain-specific kinase; EPH homology kinase 1; EHK-1; EPH-like kinase 7; EK7; hEK7
Observed Band	
Cell Pathway	Cell membrane ; Single-pass type I membrane protein . Cell projection, axon . Cell projection, dendrite .
Tissue Specificity	Almost exclusively expressed in the nervous system in cortical neurons, cerebellar Purkinje cells and pyramidal neurons within the cortex and hippocampus. Display an increasing gradient of expression from the forebrain to hindbrain and spinal cord.
Function	alternative products: Additional isoforms seem to exist, catalytic activity: ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate., function: Receptor for members of the ephrin-A family. Binds to ephrin-A1, -A2, -A3, -A4 and -A5., similarity: Belongs to the protein kinase superfamily. Tyr protein kinase family., similarity: Belongs to the protein kinase superfamily. Tyr protein kinase family. Ephrin receptor subfamily., similarity: Contains 1 protein kinase domain., similarity: Contains 1 SAM (sterile alpha motif) domain., similarity: Contains 2 fibronectin type-III domains., tissue specificity: Almost exclusively expressed in the nervous system.,
Background	This gene belongs to the ephrin receptor subfamily of the protein-tyrosine kinase family. EPH and EPH-related receptors have been implicated in mediating developmental events, particularly in the nervous system. Receptors in the EPH subfamily typically have a single kinase domain and an extracellular region



containing a Cys-rich domain and 2 fibronectin type III repeats. The ephrin receptors are divided into 2 groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands. Alternatively spliced transcript variants encoding different isoforms have been described. [provided by RefSeq, Aug 2013],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis using EphA5 Monoclonal Antibody against truncated EPHA5-His recombinant protein (1) and truncated EPHA5(aa620-774)-hIgGFc transfected CHO-K1 cell lysate(2).