

🔇 Tel: 400-999-8863 💌 Email:UpingBio@163.com

Website: www.upingBio.com

EPHA3 (Phospho Tyr602) rabbit pAb

Catalog No	YP-Ab-13130
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB; ELISA
Gene Name	EPHA3 ETK ETK1 HEK TYRO4
Protein Name	EPHA3 (Phospho Tyr602)
Immunogen	Synthesized peptide derived from human EPHA3 (Phospho Tyr602)
Specificity	This antibody detects endogenous levels of Human,Mouse,Rat EPHA3 (Phospho Tyr602)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:1000-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Ephrin type-A receptor 3 (EC 2.7.10.1;EPH-like kinase 4;EK4;hEK4;HEK;Human embryo kinase;Tyrosine-protein kinase TYRO4;Tyrosine-protein kinase receptor ETK1;Eph-like tyrosine kinase 1)
Observed Band	130kD
Cell Pathway	[Isoform 1]: Cell membrane ; Single-pass type I membrane protein .; [Isoform 2]: Secreted .
Tissue Specificity	Widely expressed. Highest level in placenta.
Function	catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate.,disease:Defects in EPHA3 may be a cause of colorectal cancer (CRC) [MIM:114500].,function:Receptor for members of the ephrin-A family. Binds to ephrin-A2, -A3, -A4 and -A5. Could play a role in lymphoid function.,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family.,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family. Ephrin receptor subfamily.,similarity:Contains 1 protein kinase domain.,similarity:Contains 1 SAM (sterile alpha motif) domain.,similarity:Contains 2 fibronectin type-III domains.,tissue specificity:Widely expressed. Highest level in placenta.,
Background	This gene belongs to the ephrin receptor subfamily of the protein-tyrosine kinase family. EPH and EPH-related receptors have been implicated in mediating



UpingBio technology Co.,Ltd

RIO		🔇 Tel: 400-999-8863 💌 Email:UpingBio@163.com 🛛 🖉 Website: www.upingBio.com
	dev sub cor rec doi Thi trai 200	velopmental events, particularly in the nervous system. Receptors in the EPH ofamily typically have a single kinase domain and an extracellular region ntaining a Cys-rich domain and 2 fibronectin type III repeats. The ephrin eptors are divided into 2 groups based on the similarity of their extracellular main sequences and their affinities for binding ephrin-A and ephrin-B ligands. Is gene encodes a protein that binds ephrin-A ligands. Two alternatively spliced inscript variants have been described for this gene. [provided by RefSeq, Jul 08],
	Av	oid repeated freezing and thawing!

matters needing attention Avoid repeated freezing and thawing:

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. Usage suggestions

Products Images

Thank you for your recent purchase