

📞 Tel: 400-999-8863 💌 Email:UpingBio@163.com

Ø Website: www.upingBio.com

## Olfactory receptor 4C6 Polyclonal Antibody

8 Mouse; A Ceptor 4C6 Im was produced against synthesized peptide derived from human range:193-242 Ceptor 4C6 Polyclonal Antibody detects endogenous levels of Ceptor 4C6 Polyclonal Antibody detects endogenous levels of Ceptor 4C6 protein. S containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Rabbit, IgG y was affinity-purified from rabbit antiserum by matography using epitope-specific immunogen. t: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: t yet tested in other applications.
A ceptor 4C6 im was produced against synthesized peptide derived from human range:193-242 ceptor 4C6 Polyclonal Antibody detects endogenous levels of ceptor 4C6 protein. S containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Rabbit,IgG y was affinity-purified from rabbit antiserum by matography using epitope-specific immunogen. tt: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA:
A ceptor 4C6 im was produced against synthesized peptide derived from human range:193-242 ceptor 4C6 Polyclonal Antibody detects endogenous levels of ceptor 4C6 protein. S containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Rabbit,IgG y was affinity-purified from rabbit antiserum by matography using epitope-specific immunogen. tt: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA:
ceptor 4C6 Im was produced against synthesized peptide derived from human range:193-242 ceptor 4C6 Polyclonal Antibody detects endogenous levels of ceptor 4C6 protein. S containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Rabbit,IgG y was affinity-purified from rabbit antiserum by matography using epitope-specific immunogen. It: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA:
im was produced against synthesized peptide derived from human range:193-242 ceptor 4C6 Polyclonal Antibody detects endogenous levels of ceptor 4C6 protein. S containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Rabbit,IgG y was affinity-purified from rabbit antiserum by matography using epitope-specific immunogen. it: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA:
im was produced against synthesized peptide derived from human range:193-242 ceptor 4C6 Polyclonal Antibody detects endogenous levels of ceptor 4C6 protein. S containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Rabbit,IgG y was affinity-purified from rabbit antiserum by matography using epitope-specific immunogen. it: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA:
range:193-242 ceptor 4C6 Polyclonal Antibody detects endogenous levels of ceptor 4C6 protein. S containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Rabbit,IgG y was affinity-purified from rabbit antiserum by matography using epitope-specific immunogen. t: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA:
ceptor 4C6 protein. S containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Rabbit,IgG y was affinity-purified from rabbit antiserum by matography using epitope-specific immunogen. it: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA:
Rabbit,IgG y was affinity-purified from rabbit antiserum by matography using epitope-specific immunogen. it: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA:
y was affinity-purified from rabbit antiserum by matography using epitope-specific immunogen. it: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA:
natography using epitope-specific immunogen. t: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA:
r
actory receptor 4C6; Olfactory receptor OR11-138
ane; Multi-pass membrane protein.
prant receptor .,similarity:Belongs to the G-protein coupled receptor 1
eptor family 4 subfamily C member 6(OR4C6) Homo sapiens ceptors interact with odorant molecules in the nose, to initiate a sponse that triggers the perception of a smell. The olfactory receptor members of a large family of G-protein-coupled receptors (GPCR) single coding-exon genes. Olfactory receptors share a brane domain structure with many neurotransmitter and hormone id are responsible for the recognition and G protein-mediated of odorant signals. The olfactory receptor gene family is the largest in The nomenclature assigned to the olfactory receptor genes and this organism is independent of other organisms. [provided by 2008],



🔇 Tel: 400-999-8863 💌 Email:UpingBio@163.com

Website: www.upingBio.com

matters needing attention

Usage suggestions

Avoid repeated freezing and thawing!

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

