

📞 Tel: 400-999-8863 💌 Email:UpingBio@163.com

Ø Website: www.upingBio.com

Olfactory receptor 4P4 Polyclonal Antibody

YP-Ab-13542 IgG Human;Rat;Mouse; WB;IF;ELISA
Human;Rat;Mouse;
WBIEFISA
OR4P4
Olfactory receptor 4P4
The antiserum was produced against synthesized peptide derived from human OR4P4. AA range:260-309
Olfactory receptor 4P4 Polyclonal Antibody detects endogenous levels of Olfactory receptor 4P4 protein.
Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Polyclonal, Rabbit,IgG
The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.
1 mg/ml
≥90%
-20°C/1 year
OR4P4; OR4P3P; Olfactory receptor 4P4; Olfactory receptor 4P3
34kD
Cell membrane; Multi-pass membrane protein.
function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,
olfactory receptor family 4 subfamily P member 4(OR4P4) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and



Avoid repeated freezing and thawing!

🔇 Tel: 400-999-8863 💌 Email:UpingBio@163.com

Website: www.upingBio.com

matters needing attention

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

