

📞 Tel: 400-999-8863 💌 Email:UpingBio@163.com

Ø Website: www.upingBio.com

Olfactory receptor 52N4 Polyclonal Antibody

Catalog No	YP-Ab-13574
lsotype	lgG
Reactivity	Human;Rat;Mouse;
Applications	WB;IF;ELISA
Gene Name	OR52N4
Protein Name	Olfactory receptor 52N4
Immunogen	The antiserum was produced against synthesized peptide derived from human OR52N4. AA range:272-321
Specificity	Olfactory receptor 52N4 Polyclonal Antibody detects endogenous levels of Olfactory receptor 52N4 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	OR52N4; Olfactory receptor 52N4; Olfactory receptor OR11-64
Observed Band	36kD
Cell Pathway	Cell membrane; Multi-pass membrane protein.
Tissue Specificity	
Function	function:Odorant receptor .,polymorphism:A stop codon at position Arg-172 in the gene coding for this protein is responsible for functional diversity thus producing a pseudogene. The stop codon is more frequent in non-Africans than in African-Americans.,similarity:Belongs to the G-protein coupled receptor 1 family.,
Background	Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],



Avoid repeated freezing and thawing!

Tel: 400-999-8863 💌 Email:UpingBio@163.com

Ø Website: www.upingBio.com

matters needing attention

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images HeI a Western Blot analysis of various cells using Olfactory (kD) receptor 52N4 Polyclonal Antibody 117-85-48-34-26-19-Western blot analysis of lysates from HeLa cells, using OR52N4 Antibody. The lane on the right is blocked with -- 117 -- 85 the synthesized peptide. -- 48 OR52N4 ---- 34 -- 26 -- 19 (kD) Western blot analysis of the lysates from COLO205 (kD) 117cells using OR52N4 antibody. 85-48-OR52N4 34-26-19-