







Olfactory receptor 5T3 Polyclonal Antibody

Catalog No YP-Ab-13607 Isotype IgG Reactivity Human;Rat;Mouse; Applications IF;ELISA Gene Name OR5T3 Protein Name Olfactory receptor 5T3 Immunogen The antiserum was produced against synthesized peptide derived from human on the protein of a smell. The olfactory receptor is near thingers the protein of a smell The rolfactory receptor arising from single coding-exon genes. Olfactory receptor-mediated of the protein of the protein on do a protein-mediated of the protein of the protein on do a protein-mediated of the protein of the protein on do protein-mediated of the protein of the protein on do protein-mediated of the protein of the protein on do protein-mediated of the protein of the protein on do protein-mediated of the protein of the protein on do protein-mediated of the protein of the protein of the pro		
Reactivity Human;Rat;Mouse; Applications IF;ELISA Gene Name OR5T3 Protein Name Olfactory receptor 5T3 Immunogen The antiserum was produced against synthesized peptide derived from hu OR5T3. AA range:86-135 Specificity Olfactory receptor 5T3 Polyclonal Antibody detects endogenous levels of Olfactory receptor 5T3 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azid Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in oth applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR5T3; Olfactory receptor 5T3; Olfactory receptor OR11-178 Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function caution: it is uncertain whether Met-1 or Met-19 is the initiator, function: Odd receptor , similarity; Belongs to the G-protein coupled receptor 1 family, olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory recroteins are members of a large family of G-protein-coupled receptors (GF arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and horma receptors and are responsible for the recognition and G protein-mediated and formar receptors and are responsible for the recognition and G protein-mediated for the recognition and G pro		Catalog No
Applications Gene Name OR5T3 Protein Name Olfactory receptor 5T3 Immunogen The antiserum was produced against synthesized peptide derived from hu OR5T3. AA range:86-135 Specificity Olfactory receptor 5T3 Polyclonal Antibody detects endogenous levels of Olfactory receptor 5T3 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azid Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in oth applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR5T3; Olfactory receptor 5T3; Olfactory receptor OR11-178 Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function caution:It is uncertain whether Met-1 or Met-19 is the initiator, function:Odd receptor .;similarity;Belongs to the G-protein coupled receptor 1 family., Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptores of arising from single coding-exon genes. Olfactory receptor share a 7-transmembrane domain structure with many neurotransmitter and hormar receptors and are responsible for the recognition and G protein-mediated		Isotype
Protein Name OR5T3 Protein Name Olfactory receptor 5T3 Immunogen The antiserum was produced against synthesized peptide derived from hu OR5T3. AA range:86-135 Specificity Olfactory receptor 5T3 Polyclonal Antibody detects endogenous levels of Olfactory receptor 5T3 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azid Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in oth applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR5T3; Olfactory receptor 5T3; Olfactory receptor OR11-178 Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function caution:It is uncertain whether Met-1 or Met-19 is the initiator, function:Odd receptor s, similarity. Belongs to the G-protein coupled receptor 1 family., Olfactory receptor family 5 subfamily T member 3(OR5T3) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors are a 7-transmembrane domain structure with many neurotransmitter and hormor receptors and are responsible for the recoption of a protein-mediated		Reactivity
Protein Name Olfactory receptor 5T3 Immunogen The antiserum was produced against synthesized peptide derived from hur OR5T3. AA range:86-135 Specificity Olfactory receptor 5T3 Polyclonal Antibody detects endogenous levels of Olfactory receptor 5T3 profein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azid Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in oth applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR5T3; Olfactory receptor 5T3; Olfactory receptor OR11-178 Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function caution:It is uncertain whether Met-1 or Met-19 is the initiator, function:Odd receptor .,simillarity:Belongs to the G-protein coupled receptor 1 family., olfactory receptor similar are receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptores interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The lofactor		Applications
Immunogen The antiserum was produced against synthesized peptide derived from hu OR5T3. AA range:86-135 Specificity Olfactory receptor 5T3 Polyclonal Antibody detects endogenous levels of Olfactory receptor 5T3 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azid Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in oth applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR5T3; Olfactory receptor 5T3; Olfactory receptor OR11-178 Observed Band Cell membrane; Multi-pass membrane protein. Tissue Specificity Function Function caution: It is uncertain whether Met-1 or Met-19 is the initiator, function: Odd receptor , similarity: Belongs to the G-protein coupled receptor 1 family. Background Olfactory receptor family 5 subfamily T member 3(OR5T3) Homo sapiens to the receptor with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptories are members of a large family of G-protein-coupled receptors (GF arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmi		Gene Name
OR5T3. AA range:86-135 Specificity Olfactory receptor 5T3 Polyclonal Antibody detects endogenous levels of Olfactory receptor 5T3 profein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azid Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in oth applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR5T3; Olfactory receptor 5T3; Olfactory receptor OR11-178 Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function caution:It is uncertain whether Met-1 or Met-19 is the initiator, function:Odd receptor .; similarity:Belongs to the G-protein coupled receptor 1 family., Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors are members of a large family of G-protein-coupled receptors (GF arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormor receptors and are responsible for the recognition and G protein-mediated		Protein Name
Olfactory receptor 5T3 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azid Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in oth applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR5T3; Olfactory receptor 5T3; Olfactory receptor OR11-178 Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function caution:It is uncertain whether Met-1 or Met-19 is the initiator, function:Odd receptor ., simillarity:Belongs to the G-protein coupled receptor 1 family. Olfactory receptor family 5 subfamily T member 3(OR5T3) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptories are members of a large family of G-protein-coupled receptors (Grarising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormor receptors and are responsible for the recognition and G protein-mediated	l from human	Immunogen
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in oth applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR5T3; Olfactory receptor 5T3; Olfactory receptor OR11-178 Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function caution:It is uncertain whether Met-1 or Met-19 is the initiator.,function:Odd receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background olfactory receptor family 5 subfamily T member 3(OR5T3) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors in a large family of G-protein-coupled receptors (Grarising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormar receptors and are responsible for the recognition and G protein-mediated	evels of	Specificity
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in oth applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR5T3; Olfactory receptor 5T3; Olfactory receptor OR11-178 Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function caution:It is uncertain whether Met-1 or Met-19 is the initiator.,function:Odd receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background olfactory receptor family 5 subfamily T member 3(OR5T3) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptoriens are members of a large family of G-protein-coupled receptors (Grarising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormor receptors and are responsible for the recognition and G protein-mediated	ium azide.	Formulation
affinity-chromatography using epitope-specific immunogen. Dilution Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in oth applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR5T3; Olfactory receptor 5T3; Olfactory receptor OR11-178 Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function caution:It is uncertain whether Met-1 or Met-19 is the initiator. function:Odd receptor .,similarity:Belongs to the G-protein coupled receptor 1 family. Background olfactory receptor family 5 subfamily T member 3(OR5T3) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptores are members of a large family of G-protein-coupled receptors (Grarising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormor receptors and are responsible for the recognition and G protein-mediated		Source
applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR5T3; Olfactory receptor 5T3; Olfactory receptor OR11-178 Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function caution:It is uncertain whether Met-1 or Met-19 is the initiator.,function:Odd receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background olfactory receptor family 5 subfamily T member 3(OR5T3) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors in are members of a large family of G-protein-coupled receptors (GF arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormor receptors and are responsible for the recognition and G protein-mediated		Purification
Purity ≥90% Storage Stability -20°C/1 year Synonyms OR5T3; Olfactory receptor 5T3; Olfactory receptor OR11-178 Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function caution:It is uncertain whether Met-1 or Met-19 is the initiator., function:Odd receptor ., similarity:Belongs to the G-protein coupled receptor 1 family., Olfactory receptor family 5 subfamily T member 3(OR5T3) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors are members of a large family of G-protein-coupled receptors (GF arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormor receptors and are responsible for the recognition and G protein-mediated	ed in other	Dilution
Synonyms OR5T3; Olfactory receptor 5T3; Olfactory receptor OR11-178 Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function caution:It is uncertain whether Met-1 or Met-19 is the initiator.,function:Odd receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background olfactory receptor family 5 subfamily T member 3(OR5T3) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors are members of a large family of G-protein-coupled receptors (GF arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormore receptors and are responsible for the recognition and G protein-mediated		Concentration
Synonyms OR5T3; Olfactory receptor 5T3; Olfactory receptor OR11-178 Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function caution:It is uncertain whether Met-1 or Met-19 is the initiator.,function:Odd receptor.,similarity:Belongs to the G-protein coupled receptor 1 family., Olfactory receptor family 5 subfamily T member 3(OR5T3) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptories are members of a large family of G-protein-coupled receptors (GF arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hornor receptors and are responsible for the recognition and G protein-mediated		Purity
Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function caution:It is uncertain whether Met-1 or Met-19 is the initiator.,function:Odd receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Olfactory receptor family 5 subfamily T member 3(OR5T3) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors are members of a large family of G-protein-coupled receptors (GF arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormoreceptors and are responsible for the recognition and G protein-mediated		Storage Stability
Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function caution:It is uncertain whether Met-1 or Met-19 is the initiator.,function:Odd receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Olfactory receptor family 5 subfamily T member 3(OR5T3) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors are members of a large family of G-protein-coupled receptors (GF arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormoreceptors and are responsible for the recognition and G protein-mediated		Synonyms
Tissue Specificity Function caution:It is uncertain whether Met-1 or Met-19 is the initiator.,function:Odd receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Olfactory receptor family 5 subfamily T member 3(OR5T3) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors are members of a large family of G-protein-coupled receptors (GF arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormore receptors and are responsible for the recognition and G protein-mediated		Observed Band
Function caution:It is uncertain whether Met-1 or Met-19 is the initiator.,function:Odd receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Olfactory receptor family 5 subfamily T member 3(OR5T3) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors are members of a large family of G-protein-coupled receptors (GF arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormore receptors and are responsible for the recognition and G protein-mediated		Cell Pathway
Background Olfactory receptor family 5 subfamily T member 3(OR5T3) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors are members of a large family of G-protein-coupled receptors (GF arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormore receptors and are responsible for the recognition and G protein-mediated		Tissue Specificity
Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors of a large family of G-protein-coupled receptors (GF arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormoreceptors and are responsible for the recognition and G protein-mediated	ction:Odorant amily.,	Function
transduction of odorant signals. The olfactory receptor gene family is the la the genome. The nomenclature assigned to the olfactory receptor genes a proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],	o initiate a actory receptor otors (GPCR) a nd hormone nediated r is the largest in	Background



UpingBio technology Co.,Ltd








matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunofluorescence analysis of MCF7 cells, using OR5T3 Antibody. The picture on the right is blocked with the synthesized peptide.