





## Olfactory receptor 8B2/3 Polyclonal Antibody

YP-Ab-13628
IgG
Human;Rat;Mouse;
WB;IF;ELISA
OR8B2/OR8B3
Olfactory receptor 8B2/3
The antiserum was produced against synthesized peptide derived from human OR8B2/8B3. AA range:199-248
Olfactory receptor 8B2/3 Polyclonal Antibody detects endogenous levels of Olfactory receptor 8B2/3 protein.
Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Polyclonal, Rabbit,IgG
The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.
1 mg/ml
≥90%
-20°C/1 year
OR8B2; Olfactory receptor 8B2; Olfactory receptor OR11-309; OR8B3; Olfactory receptor 8B3; Olfactory receptor OR11-311
35kD
Cell membrane; Multi-pass membrane protein.
function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,
olfactory receptor family 8 subfamily B member 2(OR8B2) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],



## UpingBio technology Co.,Ltd

C Tel: 400-999-8863 🛎 Email:UpingBio@163.com






matters needing attention

Avoid repeated freezing and thawing!


**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western blot analysis of lysates from Jurkat and HT-29 cells, using OR8B2/8B3 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from K562 cells using OR8B2/B3 antibody.