



Parathyroid hormone/parathyroid hormone-related peptide receptor Polyclonal Antibody

Catalog No	YP-Ab-13756
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;IHC;IF;ELISA
Gene Name	PTH1R PTHR PTHR1
Protein Name	Parathyroid hormone/parathyroid hormone-related peptide receptor
Immunogen	Synthetic peptide from human protein at AA range: 46-122
Specificity	The antibody detects endogenous Parathyroid hormone/parathyroid hormone-related peptide receptor
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000,IHC-p 1:500-200, ELISA 1:10000-20000. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Parathyroid hormone/parathyroid hormone-related peptide receptor (PTH/PTHrP type I receptor;PTH/PTHr receptor;Parathyroid hormone 1 receptor;PTH1 receptor)
Observed Band	66kD
Cell Pathway	Cell membrane ; Multi-pass membrane protein .
Tissue Specificity	Expressed in most tissues. Most abundant in kidney, bone and liver.
Function	disease:Defects in PTH1R are a cause of primary failure of tooth eruption (PFE) [MIM:125350]. PFE is a rare condition that has high penetrance and variable expressivity and in which tooth retention occurs without evidence of any obvious mechanical interference. Instead, malfunction of the eruptive mechanism itself appears to cause nonankylosed permanent teeth to fail to erupt, although the eruption pathway has been cleared by bone resorption..disease:Defects in PTH1R are the cause of chondrodysplasia Blomstrand type (BOCD) [MIM:215045]. BOCD is a severe skeletal dysplasia..disease:Defects in PTH1R are the cause of Eiken syndrome [MIM:600002]; also called Eiken skeletal dysplasia or bone modeling defect of hands and feet. Eiken syndrome is a rare familial autosomal recessive skeletal dysplasia. It is characterized by multiple

epiphyseal dysplasia, with extremely retarded ossification, pri

Background

The protein encoded by this gene is a member of the G-protein coupled receptor family 2. This protein is a receptor for parathyroid hormone (PTH) and for parathyroid hormone-like hormone (PTHrP). The activity of this receptor is mediated by G proteins which activate adenylyl cyclase and also a phosphatidylinositol-calcium second messenger system. Defects in this receptor are known to be the cause of Jansen's metaphyseal chondrodysplasia (JMC), chondrodysplasia Blomstrand type (BOCD), as well as enchondromatosis. Two transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, May 2010],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




Products Images



Western blot analysis of MCF7 Cell Lysate, antibody was diluted at 1:1000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-kidney, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-kidney, antibody was diluted at 1:200