



# C1QBP Polyclonal Antibody

<b>Catalog No</b>	YP-Ab-13872
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat
<b>Applications</b>	WB;IHC;IF;ELISA
<b>Gene Name</b>	C1QBP
<b>Protein Name</b>	Complement component 1 Q subcomponent-binding protein mitochondrial
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human C1QBP. AA range:61-110
<b>Specificity</b>	C1QBP Polyclonal Antibody detects endogenous levels of C1QBP protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/40000.. IF 1:50-200
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	C1QBP; GC1QBP; HABP1; SF2P32; Complement component 1 Q subcomponent-binding protein; mitochondrial; GC1q-R protein; Glycoprotein gC1qBP; C1qBP; Hyaluronan-binding protein 1; Mitochondrial matrix protein p32; p33
<b>Observed Band</b>	32kD
<b>Cell Pathway</b>	Mitochondrion matrix . Nucleus . Nucleus, nucleolus . Cell membrane ; Peripheral membrane protein ; Extracellular side. Secreted. Cytoplasm . Seems to be predominantly localized to mitochondria. Secreted by activated lymphocytes. Localizes to the nucleolus when coexpressed with POLGARF (PubMed:32958672). Interaction with POLGARF is likely to result in prevention of C1QBP maturation and redirection from mitochondria to nucleoli (PubMed:32958672). .
<b>Tissue Specificity</b>	Expressed on cell surface of peripheral blood cells (at protein level); Surface expression is reported for macrophages and monocyte-derived dendritic cells.
<b>Function</b>	caution:Was originally (PubMed:1830244 and PubMed:8262387) thought to be a pre-mRNA splicing factor SF2 p32 subunit and to play a role in preventing exon skipping, ensuring the accuracy of splicing and regulating alternative splicing.,function:Not known. Binds to the globular "heads" of C1Q thus inhibiting C1 activation.,similarity:Belongs to the MAM33 family.,subunit:Interacts with Rubella virus capsid protein.,



## Background

The human complement subcomponent C1q associates with C1r and C1s in order to yield the first component of the serum complement system. The protein encoded by this gene is known to bind to the globular heads of C1q molecules and inhibit C1 activation. This protein has also been identified as the p32 subunit of pre-mRNA splicing factor SF2, as well as a hyaluronic acid-binding protein. [provided by RefSeq, Jul 2008],

## matters needing attention

Avoid repeated freezing and thawing!


## Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


## Products Images



Western Blot analysis of various cells using C1QBP Polyclonal Antibody diluted at 1:1000



Immunohistochemistry analysis of paraffin-embedded human tonsil tissue, using C1QBP Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from Jurkat, HUVEC, and K562 cells, using C1QBP Antibody. The lane on the right is blocked with the synthesized peptide.