

n 🔎 WebsIte: www.upingBio.com

## TH Polyclonal Antibody

Catalog No	YP-Ab-02795
Isotype	lgG
Reactivity	Human;Mouse;Rat;Monkey
Applications	WB;IHC-p;IF(paraffin section);ELISA
Gene Name	TH
Protein Name	Tyrosine 3-monooxygenase
Immunogen	The antiserum was produced against synthesized peptide derived from human Tyrosine Hydroxylase. AA range:1-50
Specificity	TH Polyclonal Antibody detects endogenous levels of TH protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Purity Storage Stability	≥90% -20°C/1 year
•	
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20°C/1 year TH; TYH; Tyrosine 3-monooxygenase; Tyrosine 3-hydroxylase; TH 60 Cytoplasm, perinuclear region . Nucleus . Cell projection, axon . Cytoplasm .
Storage Stability Synonyms Observed Band	<ul> <li>-20°C/1 year</li> <li>TH; TYH; Tyrosine 3-monooxygenase; Tyrosine 3-hydroxylase; TH</li> <li>60</li> <li>Cytoplasm, perinuclear region . Nucleus . Cell projection, axon . Cytoplasm . Cytoplasmic vesicle, secretory vesicle, synaptic vesicle . When phosphorylated at Ser-19 shows a nuclear distribution and when phosphorylated at Ser-31 as well at Ser-40 shows a cytosolic distribution (By similarity). Expressed in dopaminergic</li> </ul>
Storage Stability Synonyms Observed Band Cell Pathway	<ul> <li>-20°C/1 year</li> <li>TH; TYH; Tyrosine 3-monooxygenase; Tyrosine 3-hydroxylase; TH</li> <li>60</li> <li>Cytoplasm, perinuclear region . Nucleus . Cell projection, axon . Cytoplasm . Cytoplasmic vesicle, secretory vesicle, synaptic vesicle . When phosphorylated at Ser-19 shows a nuclear distribution and when phosphorylated at Ser-31 as well at Ser-40 shows a cytosolic distribution (By similarity). Expressed in dopaminergic axons and axon terminals</li> </ul>
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	<ul> <li>-20°C/1 year</li> <li>TH; TYH; Tyrosine 3-monooxygenase; Tyrosine 3-hydroxylase; TH</li> <li>60</li> <li>Cytoplasm, perinuclear region . Nucleus . Cell projection, axon . Cytoplasm . Cytoplasmic vesicle, secretory vesicle, synaptic vesicle . When phosphorylated at Ser-19 shows a nuclear distribution and when phosphorylated at Ser-31 as well at Ser-40 shows a cytosolic distribution (By similarity). Expressed in dopaminergic axons and axon terminals</li> </ul>
Storage StabilitySynonymsObserved BandCell PathwayTissue SpecificityFunction	<ul> <li>-20°C/1 year</li> <li>TH; TYH; Tyrosine 3-monooxygenase; Tyrosine 3-hydroxylase; TH</li> <li>60</li> <li>Cytoplasm, perinuclear region . Nucleus . Cell projection, axon . Cytoplasm . Cytoplasmic vesicle, secretory vesicle, synaptic vesicle . When phosphorylated at Ser-19 shows a nuclear distribution and when phosphorylated at Ser-31 as well at Ser-40 shows a cytosolic distribution (By similarity). Expressed in dopaminergic axons and axon terminals</li> </ul>



🔇 Tel: 400-999-8863 📼 Email:3631691544@qq.com

Website: www.upingBio.com

## **Products Images**