

Website: www.upingBio.com

DDX5 Polyclonal Antibody

Catalog No	YP-Ab-07667
lsotype	lgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	DDX5 G17P1 HELR HLR1
Protein Name	Probable ATP-dependent RNA helicase DDX5 (EC 3.6.4.13) (DEAD box protein 5) (RNA helicase p68)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	DDX5 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
	-
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year 67 Nucleus . Nucleus, nucleolus . Cytoplasm . During the G0 phase, predominantly located in the nucleus. Cytoplasmic levels increase during the G1/S phase. During the M phase, located at the vicinity of the condensed chromosomes. At
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year 67 Nucleus . Nucleus, nucleolus . Cytoplasm . During the G0 phase, predominantly located in the nucleus. Cytoplasmic levels increase during the G1/S phase. During the M phase, located at the vicinity of the condensed chromosomes. At G1, localizes in the cytoplasm.
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year 67 Nucleus . Nucleus, nucleolus . Cytoplasm . During the G0 phase, predominantly located in the nucleus. Cytoplasmic levels increase during the G1/S phase. During the M phase, located at the vicinity of the condensed chromosomes. At G1, localizes in the cytoplasm.
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year 67 Nucleus . Nucleus, nucleolus . Cytoplasm . During the G0 phase, predominantly located in the nucleus. Cytoplasmic levels increase during the G1/S phase. During the M phase, located at the vicinity of the condensed chromosomes. At G1, localizes in the cytoplasm



🔇 Tel: 400-999-8863 📼 Email:3631691544@qq.com

Website: www.upingBio.com

Products Images