





DIAP2 Polyclonal Antibody

Catalog No	YP-Ab-07773
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	DIAPH2 DIA
Protein Name	Protein diaphanous homolog 2 (Diaphanous-related formin-2) (DRF2)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	DIAP2 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
	>000/
Purity	≥90%
Purity Storage Stability	≥90% -20°C/1 year
Storage Stability	
Storage Stability Synonyms	-20°C/1 year
Storage Stability Synonyms Observed Band	-20°C/1 year 121 [Isoform 3]: Cytoplasm, cytosol. Early endosome. Isoform 3 is cytosolic but when
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year 121 [Isoform 3]: Cytoplasm, cytosol. Early endosome. Isoform 3 is cytosolic but when coexpressed with RHOD, the 2 proteins colocalize to early endosomes. Expressed in testis, ovary, small intestine, prostate, lung, liver, kidney and
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year 121 [Isoform 3]: Cytoplasm, cytosol. Early endosome. Isoform 3 is cytosolic but when coexpressed with RHOD, the 2 proteins colocalize to early endosomes. Expressed in testis, ovary, small intestine, prostate, lung, liver, kidney and
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	-20°C/1 year 121 [Isoform 3]: Cytoplasm, cytosol. Early endosome. Isoform 3 is cytosolic but when coexpressed with RHOD, the 2 proteins colocalize to early endosomes. Expressed in testis, ovary, small intestine, prostate, lung, liver, kidney and
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	-20°C/1 year 121 [Isoform 3]: Cytoplasm, cytosol. Early endosome. Isoform 3 is cytosolic but when coexpressed with RHOD, the 2 proteins colocalize to early endosomes. Expressed in testis, ovary, small intestine, prostate, lung, liver, kidney and leukocytes.



UpingBio technology Co.,Ltd

(Tel: 400-999-8863 **(** Email:3631691544@qq.com

Website:	www.upingBio.com	

Products Images	