





## CD79a (PT1848) mouse mAb

| Catalog No   | YP-Ab-15382   |
|--|---|
| Isotype  | IgG   |
| Reactivity   | Human   |
| Applications   | IHC-p;IF(paraffin section)  |
| Gene Name  | CD79A IGA MB1   |
| Protein Name   | B-cell antigen receptor complex-associated protein alpha chain (Ig-alpha) (MB-1 membrane glycoprotein) (Membrane-bound immunoglobulin-associated protein) (Surface IgM-associated protein) (CD antigen CD79a)   |
| Immunogen  | Synthesized peptide derived from human CD79a  |
| Specificity  | This antibody detects endogenous levels of human CD79a. Heat-induced epitope retrieval (HIER) Citrate buffer of pH6.0 was highly recommended as antigen repair method in paraffin section   |
| Formulation  | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| Source   | Monoclonal, Mouse/IgG2a, Kappa  |
| Purification   | The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.  |
| Dilution   | IHC-p 1:100-500   |
| Concentration  | 1 mg/ml   |
| Purity   | ≥90%  |
|  |   |
| Storage Stability  | -20°C/1 year  |
| Storage Stability Synonyms   | -20°C/1 year  |
|  | -20°C/1 year  |
| Synonyms   | -20°C/1 year  Cell membrane; Single-pass type I membrane protein. Following antigen binding, the BCR has been shown to translocate from detergent-soluble regions of the cell membrane to lipid rafts although signal transduction through the complex can also occur outside lipid rafts |
| Synonyms Observed Band   | Cell membrane; Single-pass type I membrane protein. Following antigen binding, the BCR has been shown to translocate from detergent-soluble regions of the cell membrane to lipid rafts although signal transduction through the complex can also   |
| Synonyms  Observed Band  Cell Pathway  | Cell membrane; Single-pass type I membrane protein. Following antigen binding, the BCR has been shown to translocate from detergent-soluble regions of the cell membrane to lipid rafts although signal transduction through the complex can also occur outside lipid rafts               |
| Synonyms  Observed Band  Cell Pathway  Tissue Specificity  | Cell membrane; Single-pass type I membrane protein. Following antigen binding, the BCR has been shown to translocate from detergent-soluble regions of the cell membrane to lipid rafts although signal transduction through the complex can also occur outside lipid rafts               |
| Synonyms  Observed Band  Cell Pathway  Tissue Specificity  Function                              | Cell membrane; Single-pass type I membrane protein. Following antigen binding, the BCR has been shown to translocate from detergent-soluble regions of the cell membrane to lipid rafts although signal transduction through the complex can also occur outside lipid rafts               |
| Synonyms  Observed Band  Cell Pathway  Tissue Specificity  Function  Background  matters needing | Cell membrane; Single-pass type I membrane protein. Following antigen binding, the BCR has been shown to translocate from detergent-soluble regions of the cell membrane to lipid rafts although signal transduction through the complex can also occur outside lipid rafts  B-cells.     |



## UpingBio technology Co.,Ltd

**(** Tel: 400-999-8863 **(** Email:3631691544@qq.com

| Website: | www.upingBio.com | 1 |  |
|----------|------------------|---|--|

| Products Images |
|-----------------|
|                 |
|                 |
|                 |