

B Webslte: www.upingBio.com

TGF β Mouse mAb(1F12)

Catalog NoYP-mAb-04780IsotypeIgGReactivityHuman; Mouse;RatApplicationsWBGene NameTGFB1 TGFBProtein NameTransforming growth factor beta-1, TGF-β1, TGF bImmunogenSynthesized peptide derived from human TGF β1 MonoclonalSpecificityThis antibody detects endogenous levels of TGF β at Human, Mouse,RatFormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.11% sodium azide.SourceMonoclonal, MousePurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-1:2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsCaspase-3 (CASP-3) (EC 3.4.22.56) (Apopain) (Cysteine protease CPP32) (CPP-32) (Protein Yama) (SREBP cleavage activity 1) (SCA-1) [Cleaved into: Caspase-3 subunit p17; Caspase-3 subunit p12]Observed Band52kDCell Pathway[Latency-associated peptide]: Secreted, extracellular space, extracellular matrix .; Transforming growth factor beta-1]: Secreted.FunctionTransforming growth factor-β (TGF-β) superfamily members are critical regulators of cell proliferation and differentiation, developmental patterning and morphogenesse, and disease pathogenesis. Many cells synthesize TfB1 and have specific receptors for it. It posifively and negatively regulates many other growth factor-β (TGF-β) superfamily members are critical regulators of cell poliferation and differentiation, developmental patterning and morphogenesse, and disease pathogenesis. Many cells synthesize TfB1 and <b< th=""><th></th><th></th></b<>		
Reactivity Human; Mouse;Rat Applications WB Gene Name TGFB1 TGFB Protein Name Transforming growth factor beta-1, TGF- β 1, TGF b Immunogen Synthesized peptide derived from human TGF- β 1 Monoclonal Specificity This antibody detects endogenous levels of TGF- β at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.11% sodium azide. Source Monoclonal, Mouse Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Caspase-3 (CASP-3) (EC 3.4.22.56) (Apopain) (Cysteine protease CPP32) (CPP-32) (Protein Yama) (SREBP cleavage activity 1) (SCA-1) [Cleaved into: Caspase-3 subunit p17; Caspase-3 subunit p12] Observed Band 52kD Cell Pathway Llatency-associated peptide]: Secreted extracellular space, extracellular matrix .; Transforming growth factor beta-1]: Secreted . Tissue Specificity Highly expressed in bone (PubMed: 1746498, PubMed: 17827158). Abundantly expressed in articular cartilage and chondrocytes and is increased in cetacarthritis (OA) (PubMed: 1746498, PubMed: 17827158). Colocalizes with ASPN in	Catalog No	YP-mAb-04780
ApplicationsWBGene NameTGFB1 TGFBProtein NameTransforming growth factor beta-1, TGF-β1, TGF bImmunogenSynthesized peptide derived from human TGF β1 MonoclonalSpecificityThis antibody detects endogenous levels of TGF β at Human, Mouse, RatFormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.11% sodium azide.SourceMonoclonal, MousePurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-1:2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsCaspase-3 (CASP-3) (EC 3.4.22.56) (Apopain) (Cysteine protease CPP32) (CPP-32) (Protein Yama) (SREBP cleavage activity 1) (SCA-1) [Cleaved into: Caspase-3 subunit p17; Caspase-3 subunit p12]Observed Band52kDCell Pathway[Latency-associated peptide]: Secreted, extracellular space, extracellular matrix . (Transforming growth factor beta-1]: Secreted . Synohed: 17827158), Abundantly expressed in articular cartilage and chondrocytes and is increased in osteoarthritis (A) (PubMed: 17827158), Colocalizes with ASPN in condrocytes within AA lesions of articular cartilage (PubMed: 17827158).FunctionTransforming growth factor -β (TGF- β) superfamily members are critical regulators of cell proliferation and differentiation, developmental patterning and morphogenesis, and cisease pathogenesis. Many cells synthesize TGFB1 and have specific receptors for it. t. tpoSitvely regulates many other have specific receptors for it. t. tpoSitvely regulates many other have specific receptors for it. t. tpoSi	lsotype	lgG
Gene Name TGFB1 TGFB Protein Name Transforming growth factor beta-1, TGF- β 1, TGF b Immunogen Synthesized peptide derived from human TGF- β 1 Monoclonal Specificity This antibody detects endogenous levels of TGF- β at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.11% sodium azide. Source Monoclonal, Mouse Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Caspase-3 (CASP-3) (EC 3.4.22.56) (Apopain) (Cysteine protease CPP32) (CPP-32) (Protein Yama) (SREBP cleavage activity 1) (SCA-1) [Cleaved into: Caspase-3 subunit p17; Caspase-3 subunit p12] Observed Band 52kD Cell Pathway [Latency-associated peptide]: Secreted, extracellular space, extracellular matrix .: [Transforming growth factor beta-1]: Secreted. Tissue Specificity Tinghy expressed in orticular cartilage and chondrocytes and is increased in orticular cartilage and chondrocytes and is increased in orticular cartilage and chondrocytes and is increased in acticular cartilage and chondrocytes and is increased in orephotMed: 17827158). Abundantity expressed in articular cartilage	Reactivity	Human; Mouse;Rat
Protein Name Transforming growth factor beta-1, TGF- β 1, TGF b Immunogen Synthesized peptide derived from human TGF β 1 Monoclonal Specificity This antibody detects endogenous levels of TGF β at Human, Mouse, Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.11% sodium azide. Source Monoclonal, Mouse Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Caspase-3 (CASP-3) (EC 3.4.22.56) (Apopain) (Cysteine protease CPP32) (CPP-32) (Protein Yama) (SREBP cleavage activity 1) (SCA-1) [Cleaved into: Caspase-3 subunit p17; Caspase-3 subunit p12] Observed Band 52kD Cell Pathway [Latency-associated peptide]: Secreted, extracellular space, extracellular matrix .: [Transforming growth factor beta-1]: Secreted . Tissue Specificity Highly expressed in bone (PubMed:11746498, PubMed:17827158). Abundantly expressed in articular cartilage and chondrocytes and is increased in oeter or pubMed:11746498, PubMed:17827158). Colocalizes with ASPN in chondrocytes within OA lesions of articular cartilage synthesize TGFB 1 and have specific receptors for it. It positively and negatively feel and have specific receptors for it. It positively and negatively synthesize TGFB 1	Applications	WB
ImmunogenSynthesized peptide derived from human TGF β 1 MonoclonalSpecificityThis antibody detects endogenous levels of TGF β at Human, Mouse,RatFormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.11% sodium azide.SourceMonoclonal, MousePurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-11:2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsCaspase-3 (CASP-3) (EC 3.4.22.56) (Apopain) (Cysteine protease CPP32) (CPP-32) (Protein Yama) (SREBP cleavage activity 1) (SCA-1) [Cleaved into: Caspase-3 subunit p17; Caspase-3 subunit p12]Observed Band52kDCell Pathway[Latency-associated peptide]: Secreted, extracellular space, extracellular matrix : Transforming growth factor beta-1]: Secreted .Tissue SpecificityHighly expressed in bone (PubMed:11746498, PubMed:17827158), Abundantly expressed in articular cartilage and chondrocytes and is increased in ostocarithrits (CA) (PubMed:17426498, PubMed:17827158), Colocalizes with ASPN in chondrocytes within OA lesions of articular cartilageFunctionTransforming growth factor-β (TGF-β) superfamily members are critical regulators of cell proliferation and differentiation, developmental patterning and morphogenesis, and disease pathogenesis. Many cells synthesize TGFB1 and have specific regulators of cell proliferation and differentiation, developmental patterning and morphogenesis, and disease pathogenesis. Many cells synthesize TGFB1 and have specific regulators of cell proliferation and differentiation, developmental patterning and morp	Gene Name	TGFB1 TGFB
Specificity This antibody detects endogenous levels of TGF β at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.11% sodium azide. Source Monoclonal, Mouse Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Caspase-3 (CASP-3) (EC 3.4.22.56) (Apopain) (Cysteine protease CPP32) (CPP-32) (Protein Yama) (SREBP cleavage activity 1) (SCA-1) [Cleaved into: caspase-3 subunit p17; Caspase-3 subunit p12] Observed Band 52kD Cell Pathway [Latency-associated peptide]: Secreted , extracellular space, extracellular matrix .: [Transforming growth factor beta-1]: Secreted . Function Highly expressed in bone (PubMed:11746498, PubMed:17827158), Abundantly expressed in articular cartilage and chondrocytes and is increased in osteoarthritis (OA) (PubMed:1746498, PubMed:17827158), Abundantly expressed in articular cartilage and chondrocytes and is increased in cetoarthritis (OA) (PubMed:1746498, PubMed:17827158), Abundantly expressed in articular cartilage and chondrocytes and is increased in cetoarthritis (OA) (PubMed:17746498, PubMed:17827158), Colcalizes with ASPN in chondrocytes within OA lesions of articular cartilage (PubMed:17827158). Function Background Transforming growth factor	Protein Name	Transforming growth factor beta-1, TGF- β 1, TGF b
FormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.11% sodium azide.SourceMonoclonal, MousePurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-1:2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsCaspase-3 (CASP-3) (EC 3.4.22.56) (Apopain) (Cysteine protease CPP32) (CPP-32) (Protein Yama) (SREBP cleavage activity 1) (SCA-1) [Cleaved into: Caspase-3 subunit p17; Caspase-3 subunit p12]Observed Band52kDCell Pathway[Latency-associated peptide]: Secreted, extracellular space, extracellular matrix . (Transforming growth factor beta-1]: Secreted .Tissue SpecificityHighly expressed in bone (PubMed:11746498, PubMed:17827158). Abundantly expressed in articular cartilage and chondrocytes and is increased in osteoarthritis (OA) (PubMed:11746498, PubMed:17827158). Colocalizes with ASPP in chondrocytes within OA lesions of articular cartilage (PubMed:17827158). Colocalizes with ASPP in chondrocytes and is increased in osteoarthritis (OA) (PubMed:11746498, PubMed:17827158). Colocalizes with ASPP in chondrocytes within OA lesions of articular cartilage (PubMed:17827158).FunctionTransforming growth factor-β (TGF-β) superfamily members are critical regulators of cell proliferation and differentiation, developmental patterning and morphogenesis, and disease pathogenesis. Many cells synthesize TGFB1 and have specific receptors for it. It positively and negatively regulates many other have specific receptors for it. It positively regulates many other	Immunogen	Synthesized peptide derived from human TGF β 1 Monoclonal
SourceMonoclonal, MousePurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-1:2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsCaspase-3 (CASP-3) (EC 3.4.22.56) (Apopain) (Cysteine protease CPP32) (CPP-32) (Protein Yama) (SREBP cleavage activity 1) (SCA-1) [Cleaved into: Caspase-3 subunit p17; Caspase-3 subunit p12]Observed Band52kDCell Pathway[Latency-associated peptide]: Secreted, extracellular space, extracellular matrix : [Transforming growth factor beta-1]: Secreted.Tissue SpecificityHighly expressed in bone (PubMed:11746498, PubMed:17827158). Abundantly expressed in articular cartilage and chondrocytes and is increased in osteoarthritis (OA) (PubMed:1746498, PubMed:17827158). Colocalizes with ASPP in chondrocytes within OA lesions of articular cartilage (PubMed:17827158).FunctionTransforming growth factor-β (TGF-β) superfamily members are critical regulators of cell proliferation and differentiation, developmental patterning and morphogenesis, and disease pathogenesis. Many cells synthesize TGFB1 and have specific receptors for t. It. positively and negatively regulates many other	Specificity	This antibody detects endogenous levels of TGF β at Human, Mouse,Rat
PurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-1:2000Concentration1 mg/mlPurity $\geq 90\%$ Storage Stability-20°C/1 yearSynonymsCaspase-3 (CASP-3) (EC 3.4.22.56) (Apopain) (Cysteine protease CPP32) (CPP-32) (Protein Yama) (SREBP cleavage activity 1) (SCA-1) [Cleaved into: Caspase-3 subunit p17; Caspase-3 subunit p12]Observed Band52kDCell Pathway[Latency-associated peptide]: Secreted, extracellular space, extracellular matrix .; Transforming growth factor beta-1]: Secreted .Tissue SpecificityHighly expressed in articular cartilage and chondrocytes and is increased in osteoarthritis (OA) (PubMed:11746498, PubMed:17827158). Abundantity expressed in articular cartilage and chondrocytes and is increased in osteoarthritis (OA) (PubMed:11746498, PubMed:17827158). Colocalizes with ASPN in chondrocytes within OA lesions of articular cartilage (PubMed:17827158).FunctionTransforming growth factor- β (TGF- β) superfamily members are critical regulators of cell proliferation and differentiation, developmental patterning and morphogenesis, and disease pathogenesis. Many cells synthesize TGFB1 and have specific receptors for it. It positively and negatively regulates many other	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.11% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Caspase-3 (CASP-3) (EC 3.4.22.56) (Apopain) (Cysteine protease CPP32) (CPP-32) (Protein Yama) (SREBP cleavage activity 1) (SCA-1) [Cleaved into: Caspase-3 subunit p17; Caspase-3 subunit p12] Observed Band 52kD Cell Pathway [Latency-associated peptide]: Secreted, extracellular space, extracellular matrix .: [Transforming growth factor beta-1]: Secreted . Tissue Specificity Highly expressed in bone (PubMed:11746498, PubMed:17827158). Abundantly expressed in articular cartilage and chondrocytes and is increased in osteoarthritis (OA) (PubMed:11746498, PubMed:17827158). Colocalizes with ASPN in chondrocytes within OA lesions of articular cartilage (PubMed:17827158). Function Transforming growth factor-β (TGF-β) superfamily members are critical regulators of cell proliferation and differentiation, developmental patterning and morphogenesis, and disease pathogenesis. Many cells synthesizer TGFB1 and have specific receptors for it. It positively and negatively regulates many other	Source	Monoclonal, Mouse
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Caspase-3 (CASP-3) (EC 3.4.22.56) (Apopain) (Cysteine protease CPP32) (CPP-32) (Protein Yama) (SREBP cleavage activity 1) (SCA-1) [Cleaved into: Caspase-3 subunit p17; Caspase-3 subunit p12] Observed Band 52kD Cell Pathway [Latency-associated peptide]: Secreted, extracellular space, extracellular matrix .; ITransforming growth factor beta-1]: Secreted . Tissue Specificity Highly expressed in bone (PubMed:11746498, PubMed:17827158). Abundantly expressed in articular cartilage and chondrocytes and is increased in costeoarthritis (OA) (PubMed:1746498, PubMed:17827158). Colocalizes with ASPN in chondrocytes within OA lesions of articular cartilage Function Transforming growth factor-β (TGF-β) superfamily members are critical regulators of cell proliferation and differentiation, developmental patterning and morphogenesis, and disease pathogenesis. Many cells synthesize TGFB1 and have specific receptors for it. It positively and negatively regulates many other	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Caspase-3 (CASP-3) (EC 3.4.22.56) (Apopain) (Cysteine protease CPP32) (CPP-32) (Protein Yama) (SREBP cleavage activity 1) (SCA-1) [Cleaved into: Caspase-3 subunit p17; Caspase-3 subunit p12] Observed Band 52kD Cell Pathway [Latency-associated peptide]: Secreted, extracellular space, extracellular matrix .; [Transforming growth factor beta-1]: Secreted . Tissue Specificity Highly expressed in bone (PubMed:11746498, PubMed:17827158). Abundantly expressed in articular cartilage and chondrocytes and is increased in osteoarthritis (OA) (PubMed:11746498, PubMed:17827158). Colocalizes with ASPN in chondrocytes within OA lesions of articular cartilage (PubMed:17827158). Function Transforming growth factor- β (TGF- β) superfamily members are critical regulators of cell proliferation and differentiation, developmental patterning and morphogenesis, and disease pathogenesis. Many cells synthesize TGFB1 and have specific receptors for it. It positively and negatively regulates many other	Dilution	WB 1:500-1:2000
Storage Stability-20°C/1 yearSynonymsCaspase-3 (CASP-3) (EC 3.4.22.56) (Apopain) (Cysteine protease CPP32) (CPP-32) (Protein Yama) (SREBP cleavage activity 1) (SCA-1) [Cleaved into: Caspase-3 subunit p17; Caspase-3 subunit p12]Observed Band52kDCell Pathway[Latency-associated peptide]: Secreted, extracellular space, extracellular matrix .; [Transforming growth factor beta-1]: Secreted .Tissue SpecificityHighly expressed in bone (PubMed:11746498, PubMed:17827158). Abundantly expressed in articular cartilage and chondrocytes and is increased in osteoarthritis (OA) (PubMed:11746498, PubMed:17827158). Colocalizes with ASPN in chondrocytes within OA lesions of articular cartilage (PubMed:17827158).FunctionTransforming growth factor- β (TGF- β) superfamily members are critical regulators of cell proliferation and differentiation, developmental patterning and morphogenesis, and disease pathogenesis. Many cells synthesize TGFB1 and have specific receptors for it. It positively and negatively regulates many other	Concentration	1 mg/ml
SynonymsCaspase-3 (CASP-3) (EC 3.4.22.56) (Apopain) (Cysteine protease CPP32) (CPP-32) (Protein Yama) (SREBP cleavage activity 1) (SCA-1) [Cleaved into: Caspase-3 subunit p17; Caspase-3 subunit p12]Observed Band52kDCell Pathway[Latency-associated peptide]: Secreted, extracellular space, extracellular matrix .; [Transforming growth factor beta-1]: Secreted .Tissue SpecificityHighly expressed in bone (PubMed:11746498, PubMed:17827158). Abundantly expressed in articular cartilage and chondrocytes and is increased in osteoarthritis (OA) (PubMed:11746498, PubMed:17827158). Colocalizes with ASPN in chondrocytes within OA lesions of articular cartilage (PubMed:17827158).FunctionBackgroundTransforming growth factor-β (TGF-β) superfamily members are critical regulators of cell proliferation and differentiation, developmental patterning and morphogenesis, and disease pathogenesis. Many cells synthesize TGFB1 and have specific receptors for it. It positively and negatively regulates many other	Purity	≥90%
(CPP-32) (Protein Yama) (SREBP cleavage activity 1) (SCA-1) [Cleaved into: Caspase-3 subunit p17; Caspase-3 subunit p12]Observed Band52kDCell Pathway[Latency-associated peptide]: Secreted, extracellular space, extracellular matrix .; [Transforming growth factor beta-1]: Secreted .Tissue SpecificityHighly expressed in bone (PubMed:11746498, PubMed:17827158). Abundantly expressed in articular cartilage and chondrocytes and is increased in osteoarthritis (OA) (PubMed:11746498, PubMed:17827158). Colocalizes with ASPN in chondrocytes within OA lesions of articular cartilage (PubMed:17827158).FunctionBackgroundTransforming growth factor- β (TGF- β) superfamily members are critical regulators of cell proliferation and differentiation, developmental patterning and morphogenesis, and disease pathogenesis. Many cells synthesize TGFB1 and have specific receptors for it. It positively and negatively regulates many other	Storage Stability	-20°C/1 year
Cell Pathway[Latency-associated peptide]: Secreted, extracellular space, extracellular matrix .; [Transforming growth factor beta-1]: Secreted .Tissue SpecificityHighly expressed in bone (PubMed:11746498, PubMed:17827158). Abundantly expressed in articular cartilage and chondrocytes and is increased in osteoarthritis (OA) (PubMed:11746498, PubMed:17827158). Colocalizes with ASPN in chondrocytes within OA lesions of articular cartilage (PubMed:17827158).FunctionEackgroundTransforming growth factor- β (TGF- β) superfamily members are critical regulators of cell proliferation and differentiation, developmental patterning and morphogenesis, and disease pathogenesis. Many cells synthesize TGFB1 and have specific receptors for it. It positively and negatively regulates many other	Synonyms	(CPP-32) (Protein Yama) (SREBP cleavage activity 1) (SCA-1) [Cleaved into:
Tissue SpecificityHighly expressed in bone (PubMed:11746498, PubMed:17827158). Abundantly expressed in articular cartilage and chondrocytes and is increased in osteoarthritis (OA) (PubMed:11746498, PubMed:17827158). Colocalizes with ASPN in chondrocytes within OA lesions of articular cartilage (PubMed:17827158).FunctionTransforming growth factor- β (TGF- β) superfamily members are critical regulators of cell proliferation and differentiation, developmental patterning and morphogenesis, and disease pathogenesis. Many cells synthesize TGFB1 and have specific receptors for it. It positively and negatively regulates many other	Observed Band	52kD
osteoarthritis (OA) (PubMed:11746498, PubMed:17827158). Colocalizes with ASPN in chondrocytes within OA lesions of articular cartilage (PubMed:17827158).FunctionTransforming growth factor- β (TGF- β) superfamily members are critical regulators of cell proliferation and differentiation, developmental patterning and morphogenesis, and disease pathogenesis. Many cells synthesize TGFB1 and have specific receptors for it. It positively and negatively regulates many other	Cell Pathway	[Latency-associated peptide]: Secreted, extracellular space, extracellular matrix .; [Transforming growth factor beta-1]: Secreted .
Background Transforming growth factor- β (TGF- β) superfamily members are critical regulators of cell proliferation and differentiation, developmental patterning and morphogenesis, and disease pathogenesis. Many cells synthesize TGFB1 and have specific receptors for it. It positively and negatively regulates many other	Tissue Specificity	osteoarthritis (OA) (PubMed:11746498, PubMed:17827158). Colocalizes with ASPN in chondrocytes within OA lesions of articular cartilage
regulators of cell proliferation and differentiation, developmental patterning and morphogenesis, and disease pathogenesis. Many cells synthesize TGFB1 and have specific receptors for it. It positively and negatively regulates many other	Function	
	Background	regulators of cell proliferation and differentiation, developmental patterning and morphogenesis, and disease pathogenesis. Many cells synthesize TGFB1 and have specific receptors for it. It positively and negatively regulates many other



UpingBio technology Co.,Ltd

S Tel: 400-999-8863 Emall: UpingBio@163.com

WebsIte: www.upingBio.com

matters needing attention

Usage suggestions

Avoid repeated freezing and thawing!

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images