




# UBE3A Mouse mAb

|                           |   |
|---------------------------|---|
| Catalog No                | YP-mAb-18621  |
| Isotype                   | IgG   |
| Reactivity                | Human,Mouse,Rat   |
| Applications              | WB  |
| Gene Name                 |   |
| Protein Name              |   |
| Immunogen                 | Recombinant fusion protein containing a sequence corresponding to amino acids 300-590 of human UBE3A (NP_570853.1)  |
| Specificity               |   |
| Formulation               |   |
| Source                    |   |
| Purification              | Affinity purification   |
| Dilution                  | WB 1:500-2000   |
| Concentration             | 1 mg/ml   |
| Purity                    | ≥90%  |
| Storage Stability         | -20°C/1 year  |
| Synonyms                  | AS; ANCR; PIX1; E6-AP; HPVE6A; EPVE6AP; UBE3A   |
| Observed Band             | 100kDa  |
| Cell Pathway              |   |
| Tissue Specificity        |   |
| Function                  |   |
| Background                | This gene encodes an E3 ubiquitin-protein ligase, part of the ubiquitin protein degradation system. This imprinted gene is maternally expressed in brain and biallelically expressed in other tissues. Maternally inherited deletion of this gene causes Angelman Syndrome, characterized by severe motor and intellectual retardation, ataxia, hypotonia, epilepsy, absence of speech, and characteristic facies. The protein also interacts with the E6 protein of human papillomavirus types 16 and 18, resulting in ubiquitination and proteolysis of tumor protein p53. Alternative splicing of this gene results in three transcript variants encoding three isoforms with different N-termini. Additional transcript variants have been described, but their full length nature has not been determined. |
| matters needing attention | Avoid repeated freezing and thawing!  |

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**

Western blot analysis of various lysates, using UBE3A Mouse pAb (A1757) at 1:500 dilution. Secondary antibody: HRP Goat Anti-Mouse IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25μ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 60s.