



IgA Mouse mAb

Catalog No	YP-mAb-18622
Isotype	IgG
Reactivity	Human
Applications	WB
Gene Name	IGHA1/IGHA2
Protein Name	Ig alpha-1 chain C region/Ig alpha-2 chain C region
Immunogen	The antiserum was produced against synthesized peptide derived from human IgA. AA range:227-276
Specificity	IgA Polyclonal Antibody detects endogenous levels of IgA protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000;
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	IGHA1 ; Ig alpha-1 chain C region ; IGH A2 ; Ig alpha-2 chain C region
Observed Band	38kD
Cell Pathway	Secreted . Cell membrane .
Tissue Specificity	Bile,Colon,Mammary gland,Stomach mucosa,
Function	Disease:A chromosomal aberration involving IGH A1 is found in multiple myeloma (MM) cell lines. Translocation t(1;14)(q21;q32) that forms a FCRL4-IGH A1 fusion protein.,Function:Ig alpha is the major immunoglobulin class in body secretions. It may serve both to defend against local infection and to prevent access of foreign antigens to the general immunologic system.,PTM:3-Hydroxykynurenine, an oxidized tryptophan metabolite that is common in biological fluids, reacts with alpha-1-microglobulin to form heterogeneous polycyclic chromophores including hydroxanthommatin. The chromophore reacts with accessible cysteines forming non-reducible thioether cross-links with Ig alpha-1 chain C region Cys-352.,similarity:Contains 3 Ig-like (immunoglobulin-like) domains.,subunit:Monomeric or polymeric.,
Background	disease:A chromosomal aberration involving IGH A1 is found in multiple myeloma (MM) cell lines. Translocation t(1;14)(q21;q32) that forms a FCRL4-IGH A1 fusion protein.,function:Ig alpha is the major immunoglobulin class in body secretions. It



may serve both to defend against local infection and to prevent access of foreign antigens to the general immunologic system.,PTM:3-Hydroxykynurenine, an oxidized tryptophan metabolite that is common in biological fluids, reacts with alpha-1-microglobulin to form heterogeneous polycyclic chromophores including hydroxanthommatin. The chromophore reacts with accessible cysteines forming non-reducible thioether cross-links with Ig alpha-1 chain C region Cys-352.,similarity:Contains 3 Ig-like (immunoglobulin-like) domains.,subunit:Monomeric or polymeric.,


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using IgA Polyclonal Antibody



Western blot analysis of lysate from HeLa cells, using IgA antibody.